

10. Numerical methods: Euler's method (Notes on Diffy Qs, 1.7)

Jiří Lebl

Oklahoma State University

The textbook: <https://www.jirka.org/diffyqs/>

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

But we want answers!

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

But we want answers!

Enter numerical methods:

Given a function $f(x, y)$ and numbers x_0, y_0 , and x_1 , get (approximately) $y(x_1)$.

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

But we want answers!

Enter numerical methods:

Given a function $f(x, y)$ and numbers x_0, y_0 , and x_1 , get (approximately) $y(x_1)$.

Not ideal, but sometimes best that we got.

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

But we want answers!

Enter numerical methods:

Given a function $f(x, y)$ and numbers x_0, y_0 , and x_1 , get (approximately) $y(x_1)$.

Not ideal, but sometimes best that we got.

- We need to know $f(x, y), x_0, y_0, x_1$ exactly, we can't have parameters.

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

But we want answers!

Enter numerical methods:

Given a function $f(x, y)$ and numbers x_0, y_0 , and x_1 , get (approximately) $y(x_1)$.

Not ideal, but sometimes best that we got.

- We need to know $f(x, y), x_0, y_0, x_1$ exactly, we can't have parameters.
- Hard to get qualitative understanding from a numerical solution.

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

But we want answers!

Enter numerical methods:

Given a function $f(x, y)$ and numbers x_0, y_0 , and x_1 , get (approximately) $y(x_1)$.

Not ideal, but sometimes best that we got.

- We need to know $f(x, y), x_0, y_0, x_1$ exactly, we can't have parameters.
- Hard to get qualitative understanding from a numerical solution.
- It is only approximate, so we'd better understand the errors.

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

But we want answers!

Enter numerical methods:

Given a function $f(x, y)$ and numbers x_0, y_0 , and x_1 , get (approximately) $y(x_1)$.

Not ideal, but sometimes best that we got.

- We need to know $f(x, y), x_0, y_0, x_1$ exactly, we can't have parameters.
- Hard to get qualitative understanding from a numerical solution.
- It is only approximate, so we'd better understand the errors.
- Must choose the right method.

It is often hard to find closed form solutions to $y' = f(x, y), \quad y(x_0) = y_0.$

But we want answers!

Enter numerical methods:

Given a function $f(x, y)$ and numbers x_0, y_0 , and x_1 , get (approximately) $y(x_1)$.

Not ideal, but sometimes best that we got.

- We need to know $f(x, y), x_0, y_0, x_1$ exactly, we can't have parameters.
- Hard to get qualitative understanding from a numerical solution.
- It is only approximate, so we'd better understand the errors.
- Must choose the right method.

We will cover the simplest method of all: Euler's method.

Euler's method for $y' = f(x, y), \quad y(x_0) = y_0.$

Euler's method for $y' = f(x, y), \quad y(x_0) = y_0.$

1. Pick a step size h .

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.

1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.

1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .
3. Follow the line of slope k for x from x_0 to $x_0 + h$.

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.

1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .
3. Follow the line of slope k for x from x_0 to $x_0 + h$.
4. That is, the new point (x_1, y_1) is $x_1 = x_0 + h$ $y_1 = y_0 + hk$.

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.

1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .
3. Follow the line of slope k for x from x_0 to $x_0 + h$.
4. That is, the new point (x_1, y_1) is $x_1 = x_0 + h$ $y_1 = y_0 + hk$.
5. Compute $k = f(x_1, y_1)$.

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.

1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .
3. Follow the line of slope k for x from x_0 to $x_0 + h$.
4. That is, the new point (x_1, y_1) is $x_1 = x_0 + h$ $y_1 = y_0 + hk$.
5. Compute $k = f(x_1, y_1)$.
6. Let $x_2 = x_1 + h$ $y_2 = y_1 + hk$.

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.

1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .
3. Follow the line of slope k for x from x_0 to $x_0 + h$.
4. That is, the new point (x_1, y_1) is $x_1 = x_0 + h$ $y_1 = y_0 + hk$.
5. Compute $k = f(x_1, y_1)$.
6. Let $x_2 = x_1 + h$ $y_2 = y_1 + hk$.
7. Compute $k = f(x_2, y_2)$.

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.

1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .
3. Follow the line of slope k for x from x_0 to $x_0 + h$.
4. That is, the new point (x_1, y_1) is $x_1 = x_0 + h$ $y_1 = y_0 + hk$.
5. Compute $k = f(x_1, y_1)$.
6. Let $x_2 = x_1 + h$ $y_2 = y_1 + hk$.
7. Compute $k = f(x_2, y_2)$.
8. Let $x_3 = x_2 + h$ $y_3 = y_2 + hk$.

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.

1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .
3. Follow the line of slope k for x from x_0 to $x_0 + h$.
4. That is, the new point (x_1, y_1) is $x_1 = x_0 + h$ $y_1 = y_0 + hk$.
5. Compute $k = f(x_1, y_1)$.
6. Let $x_2 = x_1 + h$ $y_2 = y_1 + hk$.
7. Compute $k = f(x_2, y_2)$.
8. Let $x_3 = x_2 + h$ $y_3 = y_2 + hk$.
9. Rinse, repeat!

Euler's method for $y' = f(x, y)$, $y(x_0) = y_0$.


1. Pick a step size h .
2. Compute the slope $k = f(x_0, y_0)$ at (x_0, y_0) .
3. Follow the line of slope k for x from x_0 to $x_0 + h$.
4. That is, the new point (x_1, y_1) is $x_1 = x_0 + h$ $y_1 = y_0 + hk$.
5. Compute $k = f(x_1, y_1)$.
6. Let $x_2 = x_1 + h$ $y_2 = y_1 + hk$.
7. Compute $k = f(x_2, y_2)$.
8. Let $x_3 = x_2 + h$ $y_3 = y_2 + hk$.
9. Rinse, repeat!

Essentially, keep repeating : $x_{i+1} = x_i + h$, $y_{i+1} = y_i + hf(x_i, y_i)$.

Example: $y' = y^2/3, \quad y(0) = 1, \quad h = 1.$

Example: $y' = y^2/3$, $y(0) = 1$, $h = 1$.


$x_0 = 0$, $y_0 = 1$



Example: $y' = y^2/3$, $y(0) = 1$, $h = 1$.

$$x_0 = 0, \quad y_0 = 1$$

$$k = f(0, 1) = 1^2/3 = 1/3$$




Example: $y' = y^2/3$, $y(0) = 1$, $h = 1$.

$$x_0 = 0, \quad y_0 = 1$$

$$k = f(0, 1) = 1^2/3 = 1/3$$

$$x_1 = x_0 + h = 0 + 1 = 1$$




Example: $y' = y^2/3$, $y(0) = 1$, $h = 1$.

$$x_0 = 0, \quad y_0 = 1$$

$$k = f(0, 1) = 1^2/3 = 1/3$$

$$x_1 = x_0 + h = 0 + 1 = 1$$

$$y_1 = y_0 + hk = 1 + 1 \cdot (1/3) = 4/3 \approx 1.333$$



Example: $y' = y^2/3$, $y(0) = 1$, $h = 1$.


$$x_0 = 0, \quad y_0 = 1$$

$$k = f(0, 1) = 1^2/3 = 1/3$$

$$x_1 = x_0 + h = 0 + 1 = 1$$

$$y_1 = y_0 + hk = 1 + 1 \cdot (1/3) = 4/3 \approx 1.333$$

$$k = f(1, 4/3) = \frac{(4/3)^2}{3} = \frac{16}{27}$$



Example: $y' = y^2/3$, $y(0) = 1$, $h = 1$.

$$x_0 = 0, \quad y_0 = 1$$


$$k = f(0, 1) = 1^2/3 = 1/3$$

$$x_1 = x_0 + h = 0 + 1 = 1$$

$$y_1 = y_0 + hk = 1 + 1 \cdot (1/3) = 4/3 \approx 1.333$$

$$k = f(1, 4/3) = \frac{(4/3)^2}{3} = \frac{16}{27}$$

$$x_2 = x_1 + h = 1 + 1 = 2$$



Example: $y' = y^2/3$, $y(0) = 1$, $h = 1$.

$$x_0 = 0, \quad y_0 = 1$$

$$k = f(0, 1) = 1^2/3 = 1/3$$


$$x_1 = x_0 + h = 0 + 1 = 1$$

$$y_1 = y_0 + hk = 1 + 1 \cdot (1/3) = 4/3 \approx 1.333$$

$$k = f(1, 4/3) = \frac{(4/3)^2}{3} = \frac{16}{27}$$

$$x_2 = x_1 + h = 1 + 1 = 2$$

$$y_2 = y_1 + hk = 4/3 + 1 \cdot (16/27) = 52/27 \approx 1.926$$



Example: $y' = y^2/3$, $y(0) = 1$, $h = 1$.

$$x_0 = 0, \quad y_0 = 1$$

$$k = f(0, 1) = 1^2/3 = 1/3$$


$$x_1 = x_0 + h = 0 + 1 = 1$$

$$y_1 = y_0 + hk = 1 + 1 \cdot (1/3) = 4/3 \approx 1.333$$


$$k = f(1, 4/3) = \frac{(4/3)^2}{3} = \frac{16}{27}$$

$$x_2 = x_1 + h = 1 + 1 = 2$$

$$y_2 = y_1 + hk = 4/3 + 1 \cdot (16/27) = 52/27 \approx 1.926$$




Let's do a few more h s.




$$h = 1$$

Let's do a few more h s.




$$h = 1/2 = 0.5$$

Let's do a few more h s.




$$h = 1/4 = 0.25$$

Let's do a few more h s.




$$h = 1/8 = 0.125$$

Let's do a few more h s.



$$h = 1/16 = 0.0625$$

Let's do a few more h s.



$$h = 1/32 = 0.03125$$

Actual solution at $x = 2$ is $y(2) = 3$.

Actual solution at $x = 2$ is $y(2) = 3$.

The absolute value of difference between the actual solution and the approximate solution is called the *error*.

Actual solution at $x = 2$ is $y(2) = 3$.

The absolute value of difference between the actual solution and the approximate solution is called the *error*.

With $h = 1$, $x_2 = 2$ so $y(2) \approx y_2 = 1.926$

Actual solution at $x = 2$ is $y(2) = 3$.

The absolute value of difference between the actual solution and the approximate solution is called the *error*.

With $h = 1$, $x_2 = 2$ so $y(2) \approx y_2 = 1.926$ Error = $|3 - y_2| \approx 1.074$

Actual solution at $x = 2$ is $y(2) = 3$.

The absolute value of difference between the actual solution and the approximate solution is called the *error*.

With $h = 1$, $x_2 = 2$ so $y(2) \approx y_2 = 1.926$ Error = $|3 - y_2| \approx 1.074$

With $h = 1/2 = 0.5$, $x_4 = 2$ so $y(2) \approx y_4 = 2.209$

Actual solution at $x = 2$ is $y(2) = 3$.

The absolute value of difference between the actual solution and the approximate solution is called the *error*.

With $h = 1$, $x_2 = 2$ so $y(2) \approx y_2 = 1.926$ Error = $|3 - y_2| \approx 1.074$

With $h = 1/2 = 0.5$, $x_4 = 2$ so $y(2) \approx y_4 = 2.209$ Error = $|3 - y_4| \approx 0.791$

Actual solution at $x = 2$ is $y(2) = 3$.

The absolute value of difference between the actual solution and the approximate solution is called the *error*.

With $h = 1$, $x_2 = 2$ so $y(2) \approx y_2 = 1.926$ Error = $|3 - y_2| \approx 1.074$

With $h = 1/2 = 0.5$, $x_4 = 2$ so $y(2) \approx y_4 = 2.209$ Error = $|3 - y_4| \approx 0.791$

h	Approximate $y(2)$	Error	$\frac{\text{Error}}{\text{Previous error}}$
1	1.92593	1.07407	
0.5	2.20861	0.79139	0.73681
0.25	2.47250	0.52751	0.66656
0.125	2.68034	0.31966	0.60599
0.0625	2.82040	0.17960	0.56184
0.03125	2.90412	0.09588	0.53385
0.015625	2.95035	0.04965	0.51779
0.0078125	2.97472	0.02528	0.50913

Actual solution at $x = 2$ is $y(2) = 3$.

The absolute value of difference between the actual solution and the approximate solution is called the *error*.

With $h = 1$, $x_2 = 2$ so $y(2) \approx y_2 = 1.926$ Error = $|3 - y_2| \approx 1.074$

With $h = 1/2 = 0.5$, $x_4 = 2$ so $y(2) \approx y_4 = 2.209$ Error = $|3 - y_4| \approx 0.791$

h	Approximate $y(2)$	Error	$\frac{\text{Error}}{\text{Previous error}}$
1	1.92593	1.07407	
0.5	2.20861	0.79139	0.73681
0.25	2.47250	0.52751	0.66656
0.125	2.68034	0.31966	0.60599
0.0625	2.82040	0.17960	0.56184
0.03125	2.90412	0.09588	0.53385
0.015625	2.95035	0.04965	0.51779
0.0078125	2.97472	0.02528	0.50913

Note: It seems like error halves when we halve h .

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

A *second order method* reduces the error by $(1/2)^2 = 1/4$ each time you halve h .

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

A *second order method* reduces the error by $(1/2)^2 = 1/4$ each time you halve h .

A *fourth order method* reduces the error by $(1/2)^4 = 1/16$ each time you halve h .

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

A *second order method* reduces the error by $(1/2)^2 = 1/4$ each time you halve h .

A *fourth order method* reduces the error by $(1/2)^4 = 1/16$ each time you halve h .

To be within 0.1 we needed to do 64 steps of Euler.

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

A *second order method* reduces the error by $(1/2)^2 = 1/4$ each time you halve h .

A *fourth order method* reduces the error by $(1/2)^4 = 1/16$ each time you halve h .

To be within 0.1 we needed to do 64 steps of Euler.

To get within 0.01, we'd need another 2–3 halvings: so 512 to 1024 steps.

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

A *second order method* reduces the error by $(1/2)^2 = 1/4$ each time you halve h .

A *fourth order method* reduces the error by $(1/2)^4 = 1/16$ each time you halve h .

To be within 0.1 we needed to do 64 steps of Euler.

To get within 0.01, we'd need another 2–3 halvings: so 512 to 1024 steps.

If starting with one step, 10 halvings $= 2^{10} = 1024$ steps.

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

A *second order method* reduces the error by $(1/2)^2 = 1/4$ each time you halve h .

A *fourth order method* reduces the error by $(1/2)^4 = 1/16$ each time you halve h .

To be within 0.1 we needed to do 64 steps of Euler.

To get within 0.01, we'd need another 2–3 halvings: so 512 to 1024 steps.

If starting with one step, 10 halvings $= 2^{10} = 1024$ steps.

A second order method requires half as many halvings for same reduction of error.

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

A *second order method* reduces the error by $(1/2)^2 = 1/4$ each time you halve h .

A *fourth order method* reduces the error by $(1/2)^4 = 1/16$ each time you halve h .

To be within 0.1 we needed to do 64 steps of Euler.

To get within 0.01, we'd need another 2–3 halvings: so 512 to 1024 steps.

If starting with one step, 10 halvings $= 2^{10} = 1024$ steps.

A second order method requires half as many halvings for same reduction of error.

5 halvings $= 2^5 = 32$ steps.

Euler's method is a *first order method*: When we halve h , error reduces by $1/2$.

A *second order method* reduces the error by $(1/2)^2 = 1/4$ each time you halve h .

A *fourth order method* reduces the error by $(1/2)^4 = 1/16$ each time you halve h .

To be within 0.1 we needed to do 64 steps of Euler.

To get within 0.01, we'd need another 2–3 halvings: so 512 to 1024 steps.

If starting with one step, 10 halvings $= 2^{10} = 1024$ steps.

A second order method requires half as many halvings for same reduction of error.

5 halvings $= 2^5 = 32$ steps.

That can be a huge difference if we need to run this many times.

In practice we don't know the error, we must estimate it.

In practice we don't know the error, we must estimate it.

By successive halvings, and assuming it halves each time we can estimate the error.

In practice we don't know the error, we must estimate it.

By successive halvings, and assuming it halves each time we can estimate the error.

Exercise: Take the last two approximate values, and (assume they are less than $y(2)$) assuming that the error halves each time, approximate $y(2)$, and hence the error. Is it close?

h	Approximate $y(2)$	Error	$\frac{\text{Error}}{\text{Previous error}}$
1	1.92593	1.07407	
0.5	2.20861	0.79139	0.73681
0.25	2.47250	0.52751	0.66656
0.125	2.68034	0.31966	0.60599
0.0625	2.82040	0.17960	0.56184
0.03125	2.90412	0.09588	0.53385
0.015625	2.95035	0.04965	0.51779
0.0078125	2.97472	0.02528	0.50913

In practice we don't know the error, we must estimate it.

By successive halvings, and assuming it halves each time we can estimate the error.

Exercise: Take the last two approximate values, and (assume they are less than $y(2)$) assuming that the error halves each time, approximate $y(2)$, and hence the error. Is it close?

h	Approximate $y(2)$	Error	$\frac{\text{Error}}{\text{Previous error}}$
1	1.92593	1.07407	
0.5	2.20861	0.79139	0.73681
0.25	2.47250	0.52751	0.66656
0.125	2.68034	0.31966	0.60599
0.0625	2.82040	0.17960	0.56184
0.03125	2.90412	0.09588	0.53385
0.015625	2.95035	0.04965	0.51779
0.0078125	2.97472	0.02528	0.50913

The trick in practice is to estimate the error so that we can pick the right h .

Keep considering $y' = y^2/3, \quad y(0) = 1.$

Keep considering $y' = y^2/3$, $y(0) = 1$.

Suppose we attempt to find $y(3)$:

h	Approximate $y(3)$
1	3.16232
0.5	4.54329
0.25	6.86079
0.125	10.80321
0.0625	17.59893
0.03125	29.46004
0.015625	50.40121
0.0078125	87.75769

Keep considering $y' = y^2/3$, $y(0) = 1$.

Suppose we attempt to find $y(3)$:

h	Approximate $y(3)$
1	3.16232
0.5	4.54329
0.25	6.86079
0.125	10.80321
0.0625	17.59893
0.03125	29.46004
0.015625	50.40121
0.0078125	87.75769

What is going on?

Keep considering $y' = y^2/3$, $y(0) = 1$.

Suppose we attempt to find $y(3)$:

h	Approximate $y(3)$
1	3.16232
0.5	4.54329
0.25	6.86079
0.125	10.80321
0.0625	17.59893
0.03125	29.46004
0.015625	50.40121
0.0078125	87.75769

What is going on? $y(3)$ does not exist,

Keep considering $y' = y^2/3$, $y(0) = 1$.

Suppose we attempt to find $y(3)$:

h	Approximate $y(3)$
1	3.16232
0.5	4.54329
0.25	6.86079
0.125	10.80321
0.0625	17.59893
0.03125	29.46004
0.015625	50.40121
0.0078125	87.75769

What is going on? $y(3)$ does not exist, $y \rightarrow \infty$ as $x \rightarrow 3$.

In real applications, methods used are similar to Euler but better (e.g., the fourth order Runge–Kutta, see next slide).

In real applications, methods used are similar to Euler but better (e.g., the fourth order Runge–Kutta, see next slide).

Choosing h is very tricky. A couple of things to consider:

In real applications, methods used are similar to Euler but better (e.g., the fourth order Runge–Kutta, see next slide).

Choosing h is very tricky. A couple of things to consider:

- Computational time: Each step takes computer time.

In real applications, methods used are similar to Euler but better (e.g., the fourth order Runge–Kutta, see next slide).

Choosing h is very tricky. A couple of things to consider:

- Computational time: Each step takes computer time.
- Roundoff errors: Computers only compute with a certain number of significant digits.

In real applications, methods used are similar to Euler but better (e.g., the fourth order Runge–Kutta, see next slide).

Choosing h is very tricky. A couple of things to consider:

- Computational time: Each step takes computer time.
- Roundoff errors: Computers only compute with a certain number of significant digits.
- Stability: Certain equations may be numerically unstable, sometimes called *stiff* problems. Halvings may not seemingly improve things.

In real applications, methods used are similar to Euler but better (e.g., the fourth order Runge–Kutta, see next slide).

Choosing h is very tricky. A couple of things to consider:

- Computational time: Each step takes computer time.
- Roundoff errors: Computers only compute with a certain number of significant digits.
- Stability: Certain equations may be numerically unstable, sometimes called *stiff* problems. Halvings may not seemingly improve things.

In worst case, the computer might be giving us bogus numbers as if they are correct.

In real applications, methods used are similar to Euler but better (e.g., the fourth order Runge–Kutta, see next slide).

Choosing h is very tricky. A couple of things to consider:

- Computational time: Each step takes computer time.
- Roundoff errors: Computers only compute with a certain number of significant digits.
- Stability: Certain equations may be numerically unstable, sometimes called *stiff* problems. Halvings may not seemingly improve things.

In worst case, the computer might be giving us bogus numbers as if they are correct.

Numerical methods are still a current research topic.

The simplest method used in practice is the *Runge–Kutta method*.

Consider $\frac{dy}{dx} = f(x, y)$, $y(x_0) = y_0$, and a step size h .

The simplest method used in practice is the *Runge–Kutta method*.

Consider $\frac{dy}{dx} = f(x, y)$, $y(x_0) = y_0$, and a step size h .

The idea is the same as Euler except each step looks like

$$k_1 = f(x_i, y_i),$$

$$k_2 = f(x_i + h/2, y_i + k_1(h/2)),$$

$$k_3 = f(x_i + h/2, y_i + k_2(h/2)),$$

$$k_4 = f(x_i + h, y_i + k_3h).$$

$$x_{i+1} = x_i + h,$$

$$y_{i+1} = y_i + \frac{k_1 + 2k_2 + 2k_3 + k_4}{6} h,$$