

BA: 1.1

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

An *ordered set* is a set S together with a relation $<$ such that

Definition

An *ordered set* is a set S together with a relation $<$ such that

- (i) (*trichotomy*) For all $x, y \in S$, exactly one of $x < y$, $x = y$, or $y < x$ holds.

Definition

An *ordered set* is a set S together with a relation $<$ such that

- (i) (*trichotomy*) For all $x, y \in S$, exactly one of $x < y$, $x = y$, or $y < x$ holds.
- (ii) (*transitivity*) If $x, y, z \in S$ are such that $x < y$ and $y < z$, then $x < z$.

Definition

An *ordered set* is a set S together with a relation $<$ such that

- (i) (*trichotomy*) For all $x, y \in S$, exactly one of $x < y$, $x = y$, or $y < x$ holds.
- (ii) (*transitivity*) If $x, y, z \in S$ are such that $x < y$ and $y < z$, then $x < z$.

We write $x \leq y$ if $x < y$ or $x = y$. We define $>$ and \geq in the obvious way.

Definition

An *ordered set* is a set S together with a relation $<$ such that

- (i) (*trichotomy*) For all $x, y \in S$, exactly one of $x < y$, $x = y$, or $y < x$ holds.
- (ii) (*transitivity*) If $x, y, z \in S$ are such that $x < y$ and $y < z$, then $x < z$.

We write $x \leq y$ if $x < y$ or $x = y$. We define $>$ and \geq in the obvious way.

Examples:

\mathbb{Z} is an ordered set by letting $x < y$ if and only if $y - x = p$ where $p \in \mathbb{N}$.

Definition

An *ordered set* is a set S together with a relation $<$ such that

- (i) (*trichotomy*) For all $x, y \in S$, exactly one of $x < y$, $x = y$, or $y < x$ holds.
- (ii) (*transitivity*) If $x, y, z \in S$ are such that $x < y$ and $y < z$, then $x < z$.

We write $x \leq y$ if $x < y$ or $x = y$. We define $>$ and \geq in the obvious way.

Examples:

\mathbb{Z} is an ordered set by letting $x < y$ if and only if $y - x = p$ where $p \in \mathbb{N}$.

\mathbb{Q} is an ordered set by letting $x < y$ if and only if $y - x = p/q$ where $p, q \in \mathbb{N}$.

Definition

An *ordered set* is a set S together with a relation $<$ such that

- (i) (*trichotomy*) For all $x, y \in S$, exactly one of $x < y$, $x = y$, or $y < x$ holds.
- (ii) (*transitivity*) If $x, y, z \in S$ are such that $x < y$ and $y < z$, then $x < z$.

We write $x \leq y$ if $x < y$ or $x = y$. We define $>$ and \geq in the obvious way.

Examples:

\mathbb{Z} is an ordered set by letting $x < y$ if and only if $y - x = p$ where $p \in \mathbb{N}$.

\mathbb{Q} is an ordered set by letting $x < y$ if and only if $y - x = p/q$ where $p, q \in \mathbb{N}$.

The set of words is an ordered set by using lexicographic ordering.

Definition

Let $E \subset S$, where S is an ordered set.

Definition

Let $E \subset S$, where S is an ordered set.

- (i) If $\exists b \in S$ such that $x \leq b$ for all $x \in E$,
then E is *bounded above* and b is an *upper bound* of E .

Definition

Let $E \subset S$, where S is an ordered set.

- (i) If $\exists b \in S$ such that $x \leq b$ for all $x \in E$,
then E is *bounded above* and b is an *upper bound* of E .
- (ii) If $\exists b \in S$ such that $x \geq b$ for all $x \in E$,
then E is *bounded below* and b is a *lower bound* of E .

Definition

Let $E \subset S$, where S is an ordered set.

- (i) If $\exists b \in S$ such that $x \leq b$ for all $x \in E$,
then E is *bounded above* and b is an *upper bound* of E .
- (ii) If $\exists b \in S$ such that $x \geq b$ for all $x \in E$,
then E is *bounded below* and b is a *lower bound* of E .
- (iii) If \exists an upper bound b_0 of E such that $b_0 \leq b$ for all upper bounds b of E , then b_0 is called the *least upper bound* or the *supremum* of E . Write

$$\sup E := b_0.$$

Definition

Let $E \subset S$, where S is an ordered set.

- (i) If $\exists b \in S$ such that $x \leq b$ for all $x \in E$,
then E is *bounded above* and b is an *upper bound* of E .
- (ii) If $\exists b \in S$ such that $x \geq b$ for all $x \in E$,
then E is *bounded below* and b is a *lower bound* of E .
- (iii) If \exists an upper bound b_0 of E such that $b_0 \leq b$ for all upper bounds b of E , then b_0 is called the *least upper bound* or the *supremum* of E . Write

$$\sup E := b_0.$$

- (iv) If \exists a lower bound b_0 of E such that $b_0 \geq b$ for all lower bounds b of E , then b_0 is called the *greatest lower bound* or the *infimum* of E . Write

$$\inf E := b_0.$$

Definition

Let $E \subset S$, where S is an ordered set.


- (i) If $\exists b \in S$ such that $x \leq b$ for all $x \in E$,
then E is *bounded above* and b is an *upper bound* of E .
- (ii) If $\exists b \in S$ such that $x \geq b$ for all $x \in E$,
then E is *bounded below* and b is a *lower bound* of E .
- (iii) If \exists an upper bound b_0 of E such that $b_0 \leq b$ for all upper bounds b of E , then b_0 is called the *least upper bound* or the *supremum* of E . Write


$$\sup E := b_0.$$

- (iv) If \exists a lower bound b_0 of E such that $b_0 \geq b$ for all lower bounds b of E , then b_0 is called the *greatest lower bound* or the *infimum* of E . Write


$$\inf E := b_0.$$

If E is bounded above and bounded below, we say that E is *bounded*.






Notation $\sup E$ and $\inf E$ is justified as the supremum (or infimum) is unique (if it exists):



Notation $\sup E$ and $\inf E$ is justified as the supremum (or infimum) is unique (if it exists):

If b and b' are suprema of E , then $b \leq b'$ and $b' \leq b$, because both b and b' are the least upper bounds, so $b = b'$.




Notation $\sup E$ and $\inf E$ is justified as the supremum (or infimum) is unique (if it exists):

If b and b' are suprema of E , then $b \leq b'$ and $b' \leq b$, because both b and b' are the least upper bounds, so $b = b'$.

Example:

$S := \{a, b, c, d, e\}$ ordered as $a < b < c < d < e$. $E := \{a, c\}$.




Notation $\sup E$ and $\inf E$ is justified as the supremum (or infimum) is unique (if it exists):

If b and b' are suprema of E , then $b \leq b'$ and $b' \leq b$, because both b and b' are the least upper bounds, so $b = b'$.

Example:

$S := \{a, b, c, d, e\}$ ordered as $a < b < c < d < e$. $E := \{a, c\}$.

c, d , and e are upper bounds of E ,



Notation $\sup E$ and $\inf E$ is justified as the supremum (or infimum) is unique (if it exists):


If b and b' are suprema of E , then $b \leq b'$ and $b' \leq b$, because both b and b' are the least upper bounds, so $b = b'$.

Example:

$S := \{a, b, c, d, e\}$ ordered as $a < b < c < d < e$. $E := \{a, c\}$.

c, d , and e are upper bounds of E ,

c is the least upper bound or supremum of E .



Notation $\sup E$ and $\inf E$ is justified as the supremum (or infimum) is unique (if it exists):

If b and b' are suprema of E , then $b \leq b'$ and $b' \leq b$, because both b and b' are the least upper bounds, so $b = b'$.


Example:

$S := \{a, b, c, d, e\}$ ordered as $a < b < c < d < e$. $E := \{a, c\}$.

c, d , and e are upper bounds of E ,

c is the least upper bound or supremum of E .

Example: $E := \{x \in \mathbb{Q} : x < 1\} \subset \mathbb{Q}$ has a least upper bound 1, but $1 \notin E$.



Notation $\sup E$ and $\inf E$ is justified as the supremum (or infimum) is unique (if it exists):

If b and b' are suprema of E , then $b \leq b'$ and $b' \leq b$, because both b and b' are the least upper bounds, so $b = b'$.

Example:

$S := \{a, b, c, d, e\}$ ordered as $a < b < c < d < e$. $E := \{a, c\}$.

c, d , and e are upper bounds of E ,

c is the least upper bound or supremum of E .

Example: $E := \{x \in \mathbb{Q} : x < 1\} \subset \mathbb{Q}$ has a least upper bound 1, but $1 \notin E$.

Example: $P := \{x \in \mathbb{Q} : x \geq 0\} \subset \mathbb{Q}$ has no upper bound.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Suppose $x \in \mathbb{Q}$ such that $x^2 = 2$.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Suppose $x \in \mathbb{Q}$ such that $x^2 = 2$.

$x = m/n$ in lowest terms.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Suppose $x \in \mathbb{Q}$ such that $x^2 = 2$.

$x = m/n$ in lowest terms.

So $(m/n)^2 = 2$ or $m^2 = 2n^2$.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Suppose $x \in \mathbb{Q}$ such that $x^2 = 2$.

$x = m/n$ in lowest terms.

So $(m/n)^2 = 2$ or $m^2 = 2n^2$.

Hence, m^2 is divisible by 2, and so m is divisible by 2.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Suppose $x \in \mathbb{Q}$ such that $x^2 = 2$.

$x = m/n$ in lowest terms.

So $(m/n)^2 = 2$ or $m^2 = 2n^2$.

Hence, m^2 is divisible by 2, and so m is divisible by 2.

Write $m = 2k$ and so $(2k)^2 = 2n^2$.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Suppose $x \in \mathbb{Q}$ such that $x^2 = 2$.

$x = m/n$ in lowest terms.

So $(m/n)^2 = 2$ or $m^2 = 2n^2$.

Hence, m^2 is divisible by 2, and so m is divisible by 2.

Write $m = 2k$ and so $(2k)^2 = 2n^2$.

Thus $2k^2 = n^2$, and hence n is divisible by 2.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Suppose $x \in \mathbb{Q}$ such that $x^2 = 2$.

$x = m/n$ in lowest terms.

So $(m/n)^2 = 2$ or $m^2 = 2n^2$.

Hence, m^2 is divisible by 2, and so m is divisible by 2.

Write $m = 2k$ and so $(2k)^2 = 2n^2$.

Thus $2k^2 = n^2$, and hence n is divisible by 2.

Contradiction as m/n is in lowest terms.

Definition

An ordered set S has the *least-upper-bound property* if every nonempty $E \subset S$ that is bounded above has a least upper bound ($\sup E$ exists in S).

Also called the *completeness property* or the *Dedekind completeness property*.

Example: \mathbb{Q} does not have the least-upper-bound property.

$\{x \in \mathbb{Q} : x^2 < 2\}$ does not have a supremum in \mathbb{Q} .

We will show later that the supremum would be $\sqrt{2}$, but $\sqrt{2} \notin \mathbb{Q}$:

Suppose $x \in \mathbb{Q}$ such that $x^2 = 2$.

$x = m/n$ in lowest terms.

So $(m/n)^2 = 2$ or $m^2 = 2n^2$.

Hence, m^2 is divisible by 2, and so m is divisible by 2.

Write $m = 2k$ and so $(2k)^2 = 2n^2$.

Thus $2k^2 = n^2$, and hence n is divisible by 2.

Contradiction as m/n is in lowest terms.

This is the main reason why analysis needs \mathbb{R} and not just \mathbb{Q} .

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

(A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

(A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.

(A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

(A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.

(A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.

(A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

- (A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.
- (A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.
- (A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- (A4) $\exists 0 \in F$ such that $0 + x = x$ for all $x \in F$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

- (A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.
- (A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.
- (A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- (A4) $\exists 0 \in F$ such that $0 + x = x$ for all $x \in F$.
- (A5) For every $x \in F$, there exists $-x \in F$ such that $x + (-x) = 0$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

- (A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.
- (A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.
- (A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- (A4) $\exists 0 \in F$ such that $0 + x = x$ for all $x \in F$.
- (A5) For every $x \in F$, there exists $-x \in F$ such that $x + (-x) = 0$.
- (M1) $x \in F$ and $y \in F \Rightarrow xy \in F$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

- (A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.
- (A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.
- (A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- (A4) $\exists 0 \in F$ such that $0 + x = x$ for all $x \in F$.
- (A5) For every $x \in F$, there exists $-x \in F$ such that $x + (-x) = 0$.
- (M1) $x \in F$ and $y \in F \Rightarrow xy \in F$.
- (M2) (*commutativity of multiplication*) $xy = yx$ for all $x, y \in F$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

- (A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.
- (A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.
- (A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- (A4) $\exists 0 \in F$ such that $0 + x = x$ for all $x \in F$.
- (A5) For every $x \in F$, there exists $-x \in F$ such that $x + (-x) = 0$.
- (M1) $x \in F$ and $y \in F \Rightarrow xy \in F$.
- (M2) (*commutativity of multiplication*) $xy = yx$ for all $x, y \in F$.
- (M3) (*associativity of multiplication*) $(xy)z = x(yz)$ for all $x, y, z \in F$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

- (A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.
- (A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.
- (A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- (A4) $\exists 0 \in F$ such that $0 + x = x$ for all $x \in F$.
- (A5) For every $x \in F$, there exists $-x \in F$ such that $x + (-x) = 0$.
- (M1) $x \in F$ and $y \in F \Rightarrow xy \in F$.
- (M2) (*commutativity of multiplication*) $xy = yx$ for all $x, y \in F$.
- (M3) (*associativity of multiplication*) $(xy)z = x(yz)$ for all $x, y, z \in F$.
- (M4) There exists $1 \in F$ (and $1 \neq 0$) such that $1x = x$ for all $x \in F$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

- (A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.
- (A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.
- (A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- (A4) $\exists 0 \in F$ such that $0 + x = x$ for all $x \in F$.
- (A5) For every $x \in F$, there exists $-x \in F$ such that $x + (-x) = 0$.
- (M1) $x \in F$ and $y \in F \Rightarrow xy \in F$.
- (M2) (*commutativity of multiplication*) $xy = yx$ for all $x, y \in F$.
- (M3) (*associativity of multiplication*) $(xy)z = x(yz)$ for all $x, y, z \in F$.
- (M4) There exists $1 \in F$ (and $1 \neq 0$) such that $1x = x$ for all $x \in F$.
- (M5) For every $x \in F$ such that $x \neq 0$ there exists $1/x \in F$ such that $x(1/x) = 1$.

Definition

A set F is called a *field* if it has two operations defined on it, addition $x + y$ and multiplication xy , and if it satisfies the following axioms:

- (A1) $x \in F$ and $y \in F \Rightarrow x + y \in F$.
- (A2) (*commutativity of addition*) $x + y = y + x$ for all $x, y \in F$.
- (A3) (*associativity of addition*) $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- (A4) $\exists 0 \in F$ such that $0 + x = x$ for all $x \in F$.
- (A5) For every $x \in F$, there exists $-x \in F$ such that $x + (-x) = 0$.
- (M1) $x \in F$ and $y \in F \Rightarrow xy \in F$.
- (M2) (*commutativity of multiplication*) $xy = yx$ for all $x, y \in F$.
- (M3) (*associativity of multiplication*) $(xy)z = x(yz)$ for all $x, y, z \in F$.
- (M4) There exists $1 \in F$ (and $1 \neq 0$) such that $1x = x$ for all $x \in F$.
- (M5) For every $x \in F$ such that $x \neq 0$ there exists $1/x \in F$ such that $x(1/x) = 1$.
- (D) (*distributive law*) $x(y + z) = xy + xz$ for all $x, y, z \in F$.

Example: \mathbb{Q} is a field.

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Definition

A field F is said to be an *ordered field* if F is also an ordered set such that

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Definition

A field F is said to be an *ordered field* if F is also an ordered set such that

- (i) For $x, y, z \in F$, $x < y$ implies $x + z < y + z$.

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Definition

A field F is said to be an *ordered field* if F is also an ordered set such that

- (i) For $x, y, z \in F$, $x < y$ implies $x + z < y + z$.
- (ii) For $x, y \in F$, $x > 0$ and $y > 0$ implies $xy > 0$.

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Definition

A field F is said to be an *ordered field* if F is also an ordered set such that

- (i) For $x, y, z \in F$, $x < y$ implies $x + z < y + z$.
- (ii) For $x, y \in F$, $x > 0$ and $y > 0$ implies $xy > 0$.

If $x > 0$, we say x is *positive*.

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Definition

A field F is said to be an *ordered field* if F is also an ordered set such that

- (i) For $x, y, z \in F$, $x < y$ implies $x + z < y + z$.
- (ii) For $x, y \in F$, $x > 0$ and $y > 0$ implies $xy > 0$.

If $x > 0$, we say x is *positive*.

If $x < 0$, we say x is *negative*.

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Definition

A field F is said to be an *ordered field* if F is also an ordered set such that

- (i) For $x, y, z \in F$, $x < y$ implies $x + z < y + z$.
- (ii) For $x, y \in F$, $x > 0$ and $y > 0$ implies $xy > 0$.

If $x > 0$, we say x is *positive*.

If $x < 0$, we say x is *negative*.

We say x is *nonnegative* if $x \geq 0$,

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Definition

A field F is said to be an *ordered field* if F is also an ordered set such that

- (i) For $x, y, z \in F$, $x < y$ implies $x + z < y + z$.
- (ii) For $x, y \in F$, $x > 0$ and $y > 0$ implies $xy > 0$.

If $x > 0$, we say x is *positive*.

If $x < 0$, we say x is *negative*.

We say x is *nonnegative* if $x \geq 0$,

and x is *nonpositive* if $x \leq 0$.

Example: \mathbb{Q} is a field.

Example: \mathbb{Z} is not a field: No $x \in \mathbb{Z}$ such that $2x = 1$, so (M5) not satisfied.

This is not an algebra class, so we'll just assume basic properties that follow directly from the axioms without proofs.

Definition

A field F is said to be an *ordered field* if F is also an ordered set such that

- (i) For $x, y, z \in F$, $x < y$ implies $x + z < y + z$.
- (ii) For $x, y \in F$, $x > 0$ and $y > 0$ implies $xy > 0$.

If $x > 0$, we say x is *positive*.

If $x < 0$, we say x is *negative*.

We say x is *nonnegative* if $x \geq 0$,

and x is *nonpositive* if $x \leq 0$.

Example: Not hard to check that \mathbb{Q} is an ordered field.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.
- (vi) If $0 < x < y$, then $x^2 < y^2$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.
- (vi) If $0 < x < y$, then $x^2 < y^2$.
- (vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.
- (vi) If $0 < x < y$, then $x^2 < y^2$.
- (vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.

Proof: (i): $x > 0$ implies $x + (-x) > 0 + (-x)$ (item (i) of definition).

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.
- (vi) If $0 < x < y$, then $x^2 < y^2$.
- (vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.

Proof: (i): $x > 0$ implies $x + (-x) > 0 + (-x)$ (item (i) of definition).

Then $0 > -x$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.
- (vi) If $0 < x < y$, then $x^2 < y^2$.
- (vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.

Proof: (i): $x > 0$ implies $x + (-x) > 0 + (-x)$ (item (i) of definition).
Then $0 > -x$. “vice versa” follows by similar calculation.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.
- (vi) If $0 < x < y$, then $x^2 < y^2$.
- (vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.

Proof: (i): $x > 0$ implies $x + (-x) > 0 + (-x)$ (item (i) of definition).
Then $0 > -x$. “vice versa” follows by similar calculation.

(ii): $y < z$ implies $0 < z - y$ (item (i) of def).

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.
- (vi) If $0 < x < y$, then $x^2 < y^2$.
- (vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.

Proof: (i): $x > 0$ implies $x + (-x) > 0 + (-x)$ (item (i) of definition).

Then $0 > -x$. “vice versa” follows by similar calculation.

(ii): $y < z$ implies $0 < z - y$ (item (i) of def).

So by item (ii) of the definition: $0 < x(z - y)$.

Proposition

Let F be an ordered field and $x, y, z, w \in F$. Then

- (i) If $x > 0$, then $-x < 0$ (and vice versa).
- (ii) If $x > 0$ and $y < z$, then $xy < xz$.
- (iii) If $x < 0$ and $y < z$, then $xy > xz$.
- (iv) If $x \neq 0$, then $x^2 > 0$. In particular, $1 > 0$.
- (v) If $0 < x < y$, then $0 < 1/y < 1/x$.
- (vi) If $0 < x < y$, then $x^2 < y^2$.
- (vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.

Proof: (i): $x > 0$ implies $x + (-x) > 0 + (-x)$ (item (i) of definition).

Then $0 > -x$. “vice versa” follows by similar calculation.

(ii): $y < z$ implies $0 < z - y$ (item (i) of def).

So by item (ii) of the definition: $0 < x(z - y)$.

Hence $0 < xz - xy$ and by item (i) again, $xy < xz$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

$x > 0$ and $-1/x > 0 \Rightarrow x(-1/x) > 0 \Rightarrow -1 > 0$, contradicting $1 > 0$ (use (i))

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

$x > 0$ and $-1/x > 0 \Rightarrow x(-1/x) > 0 \Rightarrow -1 > 0$, contradicting $1 > 0$ (use (i))

So $1/x > 0$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

$x > 0$ and $-1/x > 0 \Rightarrow x(-1/x) > 0 \Rightarrow -1 > 0$, contradicting $1 > 0$ (use (i))

So $1/x > 0$. Similarly, $1/y > 0$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

$x > 0$ and $-1/x > 0 \Rightarrow x(-1/x) > 0 \Rightarrow -1 > 0$, contradicting $1 > 0$ (use (i))

So $1/x > 0$. Similarly, $1/y > 0$.

Thus $(1/x)(1/y) > 0$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

$x > 0$ and $-1/x > 0 \Rightarrow x(-1/x) > 0 \Rightarrow -1 > 0$, contradicting $1 > 0$ (use (i))

So $1/x > 0$. Similarly, $1/y > 0$.

Thus $(1/x)(1/y) > 0$.

By (ii) $(1/x)(1/y)x < (1/x)(1/y)y$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

$x > 0$ and $-1/x > 0 \Rightarrow x(-1/x) > 0 \Rightarrow -1 > 0$, contradicting $1 > 0$ (use (i))

So $1/x > 0$. Similarly, $1/y > 0$.

Thus $(1/x)(1/y) > 0$.

By (ii) $(1/x)(1/y)x < (1/x)(1/y)y$.

So $1/y < 1/x$.

“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x$, $z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

$x > 0$ and $-1/x > 0 \Rightarrow x(-1/x) > 0 \Rightarrow -1 > 0$, contradicting $1 > 0$ (use (i))

So $1/x > 0$. Similarly, $1/y > 0$.

Thus $(1/x)(1/y) > 0$.

By (ii) $(1/x)(1/y)x < (1/x)(1/y)y$.

So $1/y < 1/x$.

“(vi) If $0 < x < y$, then $x^2 < y^2$.”

“(vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.”

left as exercises.



“(iii) If $x < 0$ and $y < z$, then $xy > xz$.” left as exercise.

“(iv) If $x \neq 0$, then $x^2 > 0$.”:

Suppose $x > 0$. By item (ii) of definition, $x^2 > 0$.

If $x < 0$, use (iii) (with $y = x, z = 0$) to get $x^2 > 0$.

“(v) If $0 < x < y$, then $0 < 1/y < 1/x$.”:

Note $1/x \neq 0$.

If $1/x < 0$, then $-1/x > 0$ by (i).

$x > 0$ and $-1/x > 0 \Rightarrow x(-1/x) > 0 \Rightarrow -1 > 0$, contradicting $1 > 0$ (use (i))

So $1/x > 0$. Similarly, $1/y > 0$.

Thus $(1/x)(1/y) > 0$.

By (ii) $(1/x)(1/y)x < (1/x)(1/y)y$.

So $1/y < 1/x$.

“(vi) If $0 < x < y$, then $x^2 < y^2$.”

“(vii) If $x \leq y$ and $z \leq w$, then $x + z \leq y + w$.”

left as exercises. □

Example: The complex numbers \mathbb{C} (numbers $x + iy$ where $x, y \in \mathbb{R}$ and $i^2 = -1$) is not an ordered field: In every ordered field $-1 < 0$.

Proposition

Let $x, y \in F$, where F is an ordered field. If $xy > 0$, then either both x and y are positive, or both are negative.

Proposition

Let $x, y \in F$, where F is an ordered field. If $xy > 0$, then either both x and y are positive, or both are negative.

Proof: We show the contrapositive: If $x = 0$ or $y = 0$, or if x and y have opposite signs, then xy is not positive.

Proposition

Let $x, y \in F$, where F is an ordered field. If $xy > 0$, then either both x and y are positive, or both are negative.

Proof: We show the contrapositive: If $x = 0$ or $y = 0$, or if x and y have opposite signs, then xy is not positive.

If $x = 0$ or $y = 0$ is zero, then $xy = 0$ and hence not positive.

Proposition

Let $x, y \in F$, where F is an ordered field. If $xy > 0$, then either both x and y are positive, or both are negative.

Proof: We show the contrapositive: If $x = 0$ or $y = 0$, or if x and y have opposite signs, then xy is not positive.

If $x = 0$ or $y = 0$ is zero, then $xy = 0$ and hence not positive.

Suppose x and y are nonzero and have opposite signs.

Proposition

Let $x, y \in F$, where F is an ordered field. If $xy > 0$, then either both x and y are positive, or both are negative.

Proof: We show the contrapositive: If $x = 0$ or $y = 0$, or if x and y have opposite signs, then xy is not positive.

If $x = 0$ or $y = 0$ is zero, then $xy = 0$ and hence not positive.

Suppose x and y are nonzero and have opposite signs.

WLOG $x > 0$ and $y < 0$.

Proposition

Let $x, y \in F$, where F is an ordered field. If $xy > 0$, then either both x and y are positive, or both are negative.

Proof: We show the contrapositive: If $x = 0$ or $y = 0$, or if x and y have opposite signs, then xy is not positive.

If $x = 0$ or $y = 0$ is zero, then $xy = 0$ and hence not positive.

Suppose x and y are nonzero and have opposite signs.

WLOG $x > 0$ and $y < 0$.

Multiply $y < 0$ by x to get $xy < 0x = 0$.



Proposition

Let F be an ordered field with the least-upper-bound property. Let $A \subset F$ be a nonempty set that is bounded below. Then $\inf A$ exists.

Proposition

Let F be an ordered field with the least-upper-bound property. Let $A \subset F$ be a nonempty set that is bounded below. Then $\inf A$ exists.

Proof: Let $B := \{-x : x \in A\}$.

Proposition

Let F be an ordered field with the least-upper-bound property. Let $A \subset F$ be a nonempty set that is bounded below. Then $\inf A$ exists.

Proof: Let $B := \{-x : x \in A\}$.

Let $b \in F$ be a lower bound for A : If $x \in A$, then $x \geq b$. In other words, $-x \leq -b$.

Proposition

Let F be an ordered field with the least-upper-bound property. Let $A \subset F$ be a nonempty set that is bounded below. Then $\inf A$ exists.

Proof: Let $B := \{-x : x \in A\}$.

Let $b \in F$ be a lower bound for A : If $x \in A$, then $x \geq b$. In other words, $-x \leq -b$.

So $-b$ is an upper bound for B .

Proposition

Let F be an ordered field with the least-upper-bound property. Let $A \subset F$ be a nonempty set that is bounded below. Then $\inf A$ exists.

Proof: Let $B := \{-x : x \in A\}$.

Let $b \in F$ be a lower bound for A : If $x \in A$, then $x \geq b$. In other words, $-x \leq -b$.

So $-b$ is an upper bound for B .

F has the least-upper-bound property $\Rightarrow c := \sup B$ exists, and $c \leq -b$.

Proposition

Let F be an ordered field with the least-upper-bound property. Let $A \subset F$ be a nonempty set that is bounded below. Then $\inf A$ exists.

Proof: Let $B := \{-x : x \in A\}$.

Let $b \in F$ be a lower bound for A : If $x \in A$, then $x \geq b$. In other words, $-x \leq -b$.

So $-b$ is an upper bound for B .

F has the least-upper-bound property $\Rightarrow c := \sup B$ exists, and $c \leq -b$.

As $y \leq c$ for all $y \in B$, then $-c \leq x$ for all $x \in A$.

Proposition

Let F be an ordered field with the least-upper-bound property. Let $A \subset F$ be a nonempty set that is bounded below. Then $\inf A$ exists.

Proof: Let $B := \{-x : x \in A\}$.

Let $b \in F$ be a lower bound for A : If $x \in A$, then $x \geq b$. In other words, $-x \leq -b$.

So $-b$ is an upper bound for B .

F has the least-upper-bound property $\Rightarrow c := \sup B$ exists, and $c \leq -b$.

As $y \leq c$ for all $y \in B$, then $-c \leq x$ for all $x \in A$.

So $-c$ is a lower bound for A .

Proposition

Let F be an ordered field with the least-upper-bound property. Let $A \subset F$ be a nonempty set that is bounded below. Then $\inf A$ exists.

Proof: Let $B := \{-x : x \in A\}$.

Let $b \in F$ be a lower bound for A : If $x \in A$, then $x \geq b$. In other words, $-x \leq -b$.

So $-b$ is an upper bound for B .

F has the least-upper-bound property $\Rightarrow c := \sup B$ exists, and $c \leq -b$.

As $y \leq c$ for all $y \in B$, then $-c \leq x$ for all $x \in A$.

So $-c$ is a lower bound for A .

As $-c \geq b$, $-c$ is the greatest lower bound of A .

