

# BA: 1.2

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

We will simply assume that real numbers exist, that is, we won't prove:

### Theorem

*There exists a unique ordered field  $\mathbb{R}$  with the least-upper-bound property such that  $\mathbb{Q} \subset \mathbb{R}$ .*

We will simply assume that real numbers exist, that is, we won't prove:

### Theorem

*There exists a unique ordered field  $\mathbb{R}$  with the least-upper-bound property such that  $\mathbb{Q} \subset \mathbb{R}$ .*

We'll simply assume simple properties that follow from the definition.

We will simply assume that real numbers exist, that is, we won't prove:

### Theorem

*There exists a unique ordered field  $\mathbb{R}$  with the least-upper-bound property such that  $\mathbb{Q} \subset \mathbb{R}$ .*

We'll simply assume simple properties that follow from the definition.

E.g.:

$1 > 0, 2 > 0, \dots, n > 0$  for all  $n \in \mathbb{N} \subset \mathbb{R}$ .

We will simply assume that real numbers exist, that is, we won't prove:

### Theorem

*There exists a unique ordered field  $\mathbb{R}$  with the least-upper-bound property such that  $\mathbb{Q} \subset \mathbb{R}$ .*

We'll simply assume simple properties that follow from the definition.

E.g.:

$1 > 0, 2 > 0, \dots, n > 0$  for all  $n \in \mathbb{N} \subset \mathbb{R}$ .

$\frac{1}{n} > 0$  for all  $n \in \mathbb{N}$ .

We will simply assume that real numbers exist, that is, we won't prove:

### Theorem

*There exists a unique ordered field  $\mathbb{R}$  with the least-upper-bound property such that  $\mathbb{Q} \subset \mathbb{R}$ .*

We'll simply assume simple properties that follow from the definition.

E.g.:

$1 > 0, 2 > 0, \dots, n > 0$  for all  $n \in \mathbb{N} \subset \mathbb{R}$ .

$\frac{1}{n} > 0$  for all  $n \in \mathbb{N}$ .

etc.

Analysis is proving inequalities.

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ ,*



Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

$\epsilon = x/2$  gives a contradiction.



Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

$\epsilon = x/2$  gives a contradiction. □

Equivalently: *If  $x \geq 0$  is such that  $x \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

$\epsilon = x/2$  gives a contradiction. □

Equivalently: *If  $x \geq 0$  is such that  $x \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

Or the very common: *If  $|x| \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

$\epsilon = x/2$  gives a contradiction. □

Equivalently: *If  $x \geq 0$  is such that  $x \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

Or the very common: *If  $|x| \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

(To prove  $x \geq 0$ , you could prove  $-x \leq 0$  with the proposition).

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

$\epsilon = x/2$  gives a contradiction. □

Equivalently: *If  $x \geq 0$  is such that  $x \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

Or the very common: *If  $|x| \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

(To prove  $x \geq 0$ , you could prove  $-x \leq 0$  with the proposition).

We'll see many variations on the above idea.



Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

$\epsilon = x/2$  gives a contradiction.



Equivalently: *If  $x \geq 0$  is such that  $x \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

Or the very common: *If  $|x| \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

(To prove  $x \geq 0$ , you could prove  $-x \leq 0$  with the proposition).

We'll see many variations on the above idea.

Generalizing the idea of the proof:

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

$\epsilon = x/2$  gives a contradiction. □

Equivalently: *If  $x \geq 0$  is such that  $x \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

Or the very common: *If  $|x| \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

(To prove  $x \geq 0$ , you could prove  $-x \leq 0$  with the proposition).

We'll see many variations on the above idea.

Generalizing the idea of the proof:

If  $a < b$  are real numbers, then there exists  $c \in \mathbb{R}$  such that  $a < c < b$ .

Analysis is proving inequalities.

Here is how an analyst proves a nonstrict inequality:

### Proposition

*If  $x \in \mathbb{R}$  is such that  $x \leq \epsilon$  for all  $\epsilon \in \mathbb{R}$  where  $\epsilon > 0$ , then  $x \leq 0$ .*

**Proof:** Either  $x \leq 0$  or  $x > 0$ .

If  $x > 0$ , then  $0 < x/2 < x$ .

$\epsilon = x/2$  gives a contradiction. □

Equivalently: *If  $x \geq 0$  is such that  $x \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

Or the very common: *If  $|x| \leq \epsilon$  for all  $\epsilon > 0$ , then  $x = 0$ .*

(To prove  $x \geq 0$ , you could prove  $-x \leq 0$  with the proposition).

We'll see many variations on the above idea.

Generalizing the idea of the proof:

If  $a < b$  are real numbers, then there exists  $c \in \mathbb{R}$  such that  $a < c < b$ .

E.g.,  $c = \frac{a+b}{2}$ .

For analysis, the least-upper-bound property is absolutely key (allows limits).

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.



For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.

$x^2 \geq 4$  when  $x \geq 2$

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.

$x^2 \geq 4$  when  $x \geq 2 \Rightarrow x < 2$  when  $x^2 < 2$  ( $x \in A$ )

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.

$x^2 \geq 4$  when  $x \geq 2 \Rightarrow x < 2$  when  $x^2 < 2$  ( $x \in A$ )  $\Rightarrow A$  is bounded above.

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.

$x^2 \geq 4$  when  $x \geq 2 \Rightarrow x < 2$  when  $x^2 < 2$  ( $x \in A$ )  $\Rightarrow A$  is bounded above.

$1 \in A$ , so  $A \neq \emptyset$ .

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.

$x^2 \geq 4$  when  $x \geq 2 \Rightarrow x < 2$  when  $x^2 < 2$  ( $x \in A$ )  $\Rightarrow A$  is bounded above.

$1 \in A$ , so  $A \neq \emptyset$ .

So  $\sup A$  exists. Let  $r := \sup A$ .

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.

$x^2 \geq 4$  when  $x \geq 2 \Rightarrow x < 2$  when  $x^2 < 2$  ( $x \in A$ )  $\Rightarrow A$  is bounded above.

$1 \in A$ , so  $A \neq \emptyset$ .

So  $\sup A$  exists. Let  $r := \sup A$ .

Immediate:  $r \geq 1 > 0$  as  $1 \in A$ .

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.

$x^2 \geq 4$  when  $x \geq 2 \Rightarrow x < 2$  when  $x^2 < 2$  ( $x \in A$ )  $\Rightarrow A$  is bounded above.

$1 \in A$ , so  $A \neq \emptyset$ .

So  $\sup A$  exists. Let  $r := \sup A$ .

Immediate:  $r \geq 1 > 0$  as  $1 \in A$ .

WTS  $r^2 = 2$ .

For analysis, the least-upper-bound property is absolutely key (allows limits).

Let's use it to show that  $\sqrt{2} \in \mathbb{R}$  exists (it does not exist in  $\mathbb{Q}$ ).

**Example:**  $\exists$  a unique positive  $r \in \mathbb{R}$  such that  $r^2 = 2$ .

**Proof:** Let  $A := \{x \in \mathbb{R} : x^2 < 2\}$ .

We claim that  $A$  is bounded above and nonempty.

$x^2 \geq 4$  when  $x \geq 2 \Rightarrow x < 2$  when  $x^2 < 2$  ( $x \in A$ )  $\Rightarrow A$  is bounded above.

$1 \in A$ , so  $A \neq \emptyset$ .

So  $\sup A$  exists. Let  $r := \sup A$ .

Immediate:  $r \geq 1 > 0$  as  $1 \in A$ .

WTS  $r^2 = 2$ .

This is analysis, we'll show  $r^2 \geq 2$  and  $r^2 \leq 2$ .



**Claim:**  $r^2 \geq 2$ .

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

Take  $h \in \mathbb{R}$  such that  $0 < h < \frac{2-s^2}{2s+1}$  and  $h < 1$ .

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

Take  $h \in \mathbb{R}$  such that  $0 < h < \frac{2-s^2}{2s+1}$  and  $h < 1$ .

$$(s + h)^2 - s^2 = h(2s + h)$$

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

Take  $h \in \mathbb{R}$  such that  $0 < h < \frac{2-s^2}{2s+1}$  and  $h < 1$ .

$$\begin{aligned}(s + h)^2 - s^2 &= h(2s + h) \\ &< h(2s + 1) \quad (\text{since } h < 1)\end{aligned}$$

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

Take  $h \in \mathbb{R}$  such that  $0 < h < \frac{2-s^2}{2s+1}$  and  $h < 1$ .

$$(s + h)^2 - s^2 = h(2s + h)$$

$$< h(2s + 1) \quad (\text{since } h < 1)$$

$$< 2 - s^2 \quad (\text{since } h < \frac{2-s^2}{2s+1})$$

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

Take  $h \in \mathbb{R}$  such that  $0 < h < \frac{2-s^2}{2s+1}$  and  $h < 1$ .

$$\begin{aligned}(s + h)^2 - s^2 &= h(2s + h) \\ &< h(2s + 1) \quad (\text{since } h < 1) \\ &< 2 - s^2 \quad (\text{since } h < \frac{2-s^2}{2s+1})\end{aligned}$$

$$\Rightarrow (s + h)^2 < 2 \quad \Rightarrow \quad s + h \in A$$



**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

Take  $h \in \mathbb{R}$  such that  $0 < h < \frac{2-s^2}{2s+1}$  and  $h < 1$ .

$$\begin{aligned}(s + h)^2 - s^2 &= h(2s + h) \\ &< h(2s + 1) \quad (\text{since } h < 1) \\ &< 2 - s^2 \quad (\text{since } h < \frac{2-s^2}{2s+1})\end{aligned}$$

$$\Rightarrow (s + h)^2 < 2 \quad \Rightarrow \quad s + h \in A$$

$$\text{but } s + h > s \text{ as } h > 0 \quad \Rightarrow \quad s < r = \sup A$$

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

Take  $h \in \mathbb{R}$  such that  $0 < h < \frac{2-s^2}{2s+1}$  and  $h < 1$ .

$$\begin{aligned}(s + h)^2 - s^2 &= h(2s + h) \\ &< h(2s + 1) \quad (\text{since } h < 1) \\ &< 2 - s^2 \quad (\text{since } h < \frac{2-s^2}{2s+1})\end{aligned}$$

$$\Rightarrow (s + h)^2 < 2 \quad \Rightarrow \quad s + h \in A$$

$$\text{but } s + h > s \text{ as } h > 0 \quad \Rightarrow \quad s < r = \sup A$$

$$s \text{ was arbitrary positive } s \text{ such that } s^2 < 2 \quad \Rightarrow \quad r^2 \geq 2.$$

**Claim:**  $r^2 \geq 2$ .

**Proof of claim:** Take  $s > 0$  such that  $s^2 < 2$ . Want  $h > 0$  such that  $(s + h)^2 < 2$ .

$$2 - s^2 > 0 \quad \Rightarrow \quad \frac{2-s^2}{2s+1} > 0.$$

Take  $h \in \mathbb{R}$  such that  $0 < h < \frac{2-s^2}{2s+1}$  and  $h < 1$ .

$$\begin{aligned}(s + h)^2 - s^2 &= h(2s + h) \\ &< h(2s + 1) \quad (\text{since } h < 1) \\ &< 2 - s^2 \quad (\text{since } h < \frac{2-s^2}{2s+1})\end{aligned}$$

$$\Rightarrow (s + h)^2 < 2 \quad \Rightarrow \quad s + h \in A$$

$$\text{but } s + h > s \text{ as } h > 0 \quad \Rightarrow \quad s < r = \sup A$$

$$s \text{ was arbitrary positive } s \text{ such that } s^2 < 2 \quad \Rightarrow \quad r^2 \geq 2.$$

Claim is proved.

**Claim:**  $r^2 \leq 2$ .

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ .

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ .    Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ .    Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \implies h = \frac{s^2 - 2}{2s} > 0$ .



**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \implies h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2-2}{2s}$ .  $s^2 - 2 > 0 \implies h = \frac{s^2-2}{2s} > 0$ .  $s - h = s - \frac{s^2-2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$s^2 - (s - h)^2 = 2sh - h^2$$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \implies h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \end{aligned}$$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2-2}{2s}$ .  $s^2 - 2 > 0 \implies h = \frac{s^2-2}{2s} > 0$ .  $s - h = s - \frac{s^2-2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$s^2 - (s - h)^2 = 2sh - h^2$$

$$< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0)$$

$$= s^2 - 2 \quad (\text{since } h = \frac{s^2-2}{2s})$$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2-2}{2s}$ .  $s^2 - 2 > 0 \implies h = \frac{s^2-2}{2s} > 0$ .  $s - h = s - \frac{s^2-2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$s^2 - (s - h)^2 = 2sh - h^2$$

$$< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0)$$

$$= s^2 - 2 \quad (\text{since } h = \frac{s^2-2}{2s})$$

$$\implies (s - h)^2 > 2$$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2-2}{2s}$ .  $s^2 - 2 > 0 \implies h = \frac{s^2-2}{2s} > 0$ .  $s - h = s - \frac{s^2-2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$s^2 - (s - h)^2 = 2sh - h^2$$

$$< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0)$$

$$= s^2 - 2 \quad (\text{since } h = \frac{s^2-2}{2s})$$

$$\implies (s - h)^2 > 2 \implies s - h \notin A.$$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2-2}{2s}$ .  $s^2 - 2 > 0 \implies h = \frac{s^2-2}{2s} > 0$ .  $s - h = s - \frac{s^2-2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2-2}{2s}) \end{aligned}$$

$$\implies (s - h)^2 > 2 \implies s - h \notin A.$$

$$\text{If } x \geq s - h \implies x^2 \geq (s - h)^2 > 2$$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2-2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2-2}{2s} > 0$ .  $s - h = s - \frac{s^2-2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2-2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

$$\text{If } x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A$$



**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2-2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2-2}{2s} > 0$ .  $s - h = s - \frac{s^2-2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2-2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2 - 2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2.$$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2 - 2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2$ . The claim is proved.

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2 - 2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2$ . The claim is proved.

$$r^2 \geq 2 \text{ and } r^2 \leq 2 \Rightarrow r^2 = 2.$$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2 - 2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2$ . The claim is proved.

$r^2 \geq 2$  and  $r^2 \leq 2 \Rightarrow r^2 = 2$ . Existence is done.

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2 - 2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2$ . The claim is proved.

$r^2 \geq 2$  and  $r^2 \leq 2 \Rightarrow r^2 = 2$ . Existence is done.

Uniqueness: Suppose  $s > 0$  such that  $s^2 = 2$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2 - 2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2$ . The claim is proved.

$r^2 \geq 2$  and  $r^2 \leq 2 \Rightarrow r^2 = 2$ . Existence is done.

Uniqueness: Suppose  $s > 0$  such that  $s^2 = 2 \Rightarrow s^2 = r^2$ .

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2 - 2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2$ . The claim is proved.

$r^2 \geq 2$  and  $r^2 \leq 2 \Rightarrow r^2 = 2$ . Existence is done.

Uniqueness: Suppose  $s > 0$  such that  $s^2 = 2 \Rightarrow s^2 = r^2$ .

$0 < s < r$  would imply  $s^2 < r^2$



**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2-2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2-2}{2s} > 0$ .  $s - h = s - \frac{s^2-2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2-2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2$ . The claim is proved.

$r^2 \geq 2$  and  $r^2 \leq 2 \Rightarrow r^2 = 2$ . Existence is done.

Uniqueness: Suppose  $s > 0$  such that  $s^2 = 2 \Rightarrow s^2 = r^2$ .

$0 < s < r$  would imply  $s^2 < r^2$   $0 < r < s$  would imply  $r^2 < s^2$

**Claim:**  $r^2 \leq 2$ .

**Proof of claim:**

Take  $s > 0$  such that  $s^2 > 2$ . Want  $h > 0$  such that  $(s - h)^2 > 2$  and  $s - h > 0$ .

Let  $h := \frac{s^2 - 2}{2s}$ .  $s^2 - 2 > 0 \Rightarrow h = \frac{s^2 - 2}{2s} > 0$ .  $s - h = s - \frac{s^2 - 2}{2s} = \frac{s}{2} + \frac{1}{s} > 0$ .

$$\begin{aligned} s^2 - (s - h)^2 &= 2sh - h^2 \\ &< 2sh \quad (\text{since } h^2 > 0 \text{ as } h \neq 0) \\ &= s^2 - 2 \quad (\text{since } h = \frac{s^2 - 2}{2s}) \end{aligned}$$

$$\Rightarrow (s - h)^2 > 2 \Rightarrow s - h \notin A.$$

If  $x \geq s - h \Rightarrow x^2 \geq (s - h)^2 > 2 \Rightarrow x \notin A \Rightarrow s - h$  is an upper bound.

$s - h < s \Rightarrow s > r = \sup A \Rightarrow r^2 \leq 2$ . The claim is proved.

$r^2 \geq 2$  and  $r^2 \leq 2 \Rightarrow r^2 = 2$ . Existence is done.

Uniqueness: Suppose  $s > 0$  such that  $s^2 = 2 \Rightarrow s^2 = r^2$ .

$0 < s < r$  would imply  $s^2 < r^2$   $0 < r < s$  would imply  $r^2 < s^2 \Rightarrow s = r$ .

□

$x^{1/n}$  exists for any  $n \in \mathbb{N}$  and all  $x > 0$  (proof harder).

$x^{1/n}$  exists for any  $n \in \mathbb{N}$  and all  $x > 0$  (proof harder).

$$\sqrt{2} \notin \mathbb{Q}$$

$x^{1/n}$  exists for any  $n \in \mathbb{N}$  and all  $x > 0$  (proof harder).

$\sqrt{2} \notin \mathbb{Q}$       so  $\mathbb{R} \setminus \mathbb{Q} \neq \emptyset$

$x^{1/n}$  exists for any  $n \in \mathbb{N}$  and all  $x > 0$  (proof harder).

$\sqrt{2} \notin \mathbb{Q}$       so  $\mathbb{R} \setminus \mathbb{Q} \neq \emptyset$

(In fact, we'll see  $\mathbb{R} \setminus \mathbb{Q}$  is much bigger than  $\mathbb{Q}$ .)

$x^{1/n}$  exists for any  $n \in \mathbb{N}$  and all  $x > 0$  (proof harder).

$\sqrt{2} \notin \mathbb{Q}$       so  $\mathbb{R} \setminus \mathbb{Q} \neq \emptyset$

(In fact, we'll see  $\mathbb{R} \setminus \mathbb{Q}$  is much bigger than  $\mathbb{Q}$ .)

$\mathbb{R} \setminus \mathbb{Q}$  is called the set of *irrational* numbers.

## Theorem

(i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .



## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i):

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ .

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ . (i) says that  $\forall$  real  $t := y/x \exists n \in \mathbb{N}$  such that  $n > t$ .

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ . (i) says that  $\forall$  real  $t := y/x \exists n \in \mathbb{N}$  such that  $n > t$ .

So (i) says that  $\mathbb{N} \subset \mathbb{R}$  is not bounded above.

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ . (i) says that  $\forall$  real  $t := y/x \exists n \in \mathbb{N}$  such that  $n > t$ .

So (i) says that  $\mathbb{N} \subset \mathbb{R}$  is not bounded above.

Suppose  $\mathbb{N}$  is bounded (for contradiction).

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ . (i) says that  $\forall$  real  $t := y/x \exists n \in \mathbb{N}$  such that  $n > t$ .

So (i) says that  $\mathbb{N} \subset \mathbb{R}$  is not bounded above.

Suppose  $\mathbb{N}$  is bounded (for contradiction).

Let  $b := \sup \mathbb{N}$ .



## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ . (i) says that  $\forall$  real  $t := y/x \exists n \in \mathbb{N}$  such that  $n > t$ .

So (i) says that  $\mathbb{N} \subset \mathbb{R}$  is not bounded above.

Suppose  $\mathbb{N}$  is bounded (for contradiction).

Let  $b := \sup \mathbb{N}$ .

$b - 1$  is not an upper bound as  $b - 1 < b$ .

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ . (i) says that  $\forall$  real  $t := y/x \exists n \in \mathbb{N}$  such that  $n > t$ .

So (i) says that  $\mathbb{N} \subset \mathbb{R}$  is not bounded above.

Suppose  $\mathbb{N}$  is bounded (for contradiction).

Let  $b := \sup \mathbb{N}$ .

$b - 1$  is not an upper bound as  $b - 1 < b$ .

$\Rightarrow \exists m \in \mathbb{N}$  such that  $m > b - 1$

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ . (i) says that  $\forall$  real  $t := y/x \exists n \in \mathbb{N}$  such that  $n > t$ .

So (i) says that  $\mathbb{N} \subset \mathbb{R}$  is not bounded above.

Suppose  $\mathbb{N}$  is bounded (for contradiction).

Let  $b := \sup \mathbb{N}$ .

$b - 1$  is not an upper bound as  $b - 1 < b$ .

$\Rightarrow \exists m \in \mathbb{N}$  such that  $m > b - 1 \Rightarrow m + 1 > b (\Rightarrow \Leftarrow)$

## Theorem

- (i) (Archimedean property) If  $x, y \in \mathbb{R}$  and  $x > 0$ , then  $\exists n \in \mathbb{N}$  such that  $nx > y$ .
- (ii) ( $\mathbb{Q}$  is dense in  $\mathbb{R}$ ) If  $x, y \in \mathbb{R}$  and  $x < y$ , then  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Remark: The two parts are actually equivalent.

**Proof:** (i): Divide by  $x$ . (i) says that  $\forall$  real  $t := y/x \exists n \in \mathbb{N}$  such that  $n > t$ .

So (i) says that  $\mathbb{N} \subset \mathbb{R}$  is not bounded above.

Suppose  $\mathbb{N}$  is bounded (for contradiction).

Let  $b := \sup \mathbb{N}$ .

$b - 1$  is not an upper bound as  $b - 1 < b$ .


$\Rightarrow \exists m \in \mathbb{N}$  such that  $m > b - 1 \Rightarrow m + 1 > b (\Rightarrow \Leftarrow)$

(i) is proved.

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .




Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .




Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .



Idea for (ii) (density of  $\mathbb{Q}$ ):


Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).





Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,


then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .




Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .



$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).


Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .



(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .


$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .



$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .


If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .



$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .


If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .



(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).


$n(y - x) > 1 \Rightarrow ny > 1 + nx$

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .



$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).


$n(y - x) > 1 \Rightarrow ny > 1 + nx \Rightarrow ny > 1 + nx \geq m$

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .



$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).

$n(y - x) > 1 \Rightarrow ny > 1 + nx \Rightarrow ny > 1 + nx \geq m \Rightarrow y > m/n$




Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .



$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).


$n(y - x) > 1 \Rightarrow ny > 1 + nx \Rightarrow ny > 1 + nx \geq m \Rightarrow y > m/n$  So  $r = m/n$  works.

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .

(ii): First assume  $x \geq 0$ .



$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).


$n(y - x) > 1 \Rightarrow ny > 1 + nx \Rightarrow ny > 1 + nx \geq m \Rightarrow y > m/n$  So  $r = m/n$  works.

Now assume  $x < 0$ .

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .



(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).


$n(y - x) > 1 \Rightarrow ny > 1 + nx \Rightarrow ny > 1 + nx \geq m \Rightarrow y > m/n$  So  $r = m/n$  works.

Now assume  $x < 0$ . If  $y > 0$ , then  $r = 0$  works.

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .



(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).

$n(y - x) > 1 \Rightarrow ny > 1 + nx \Rightarrow ny > 1 + nx \geq m \Rightarrow y > m/n$  So  $r = m/n$  works.


Now assume  $x < 0$ . If  $y > 0$ , then  $r = 0$  works.

If  $y \leq 0$ , then  $0 \leq -y < -x$ .

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .



(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).

$n(y - x) > 1 \Rightarrow ny > 1 + nx \Rightarrow ny > 1 + nx \geq m \Rightarrow y > m/n$  So  $r = m/n$  works.


Now assume  $x < 0$ . If  $y > 0$ , then  $r = 0$  works.

If  $y \leq 0$ , then  $0 \leq -y < -x$ . Find  $q \in \mathbb{Q}$  s.t.  $-y < q < -x$ .

Idea for (ii) (density of  $\mathbb{Q}$ ):

Find  $n$  such that  $y - x > 1/n$ ,

then find least  $m$  such that  $m/n > x$ .



(ii): First assume  $x \geq 0$ .

$y - x > 0$  (using (i))  $\Rightarrow \exists n \in \mathbb{N}$  such that  $n(y - x) > 1$  or  $y - x > 1/n$ .

$A := \{k \in \mathbb{N} : k > nx\}$  is nonempty by (i).

Well ordering property  $\Rightarrow \exists$  a least element  $m$  of  $A$ .

$m \in A \Rightarrow m > nx \Rightarrow x < m/n$ .

$m$  is the least element  $\Rightarrow m - 1 \notin A$ .

If  $m > 1$ , then  $m - 1 \in \mathbb{N}$  and  $m - 1 \notin A$  so  $m - 1 \leq nx$ .

If  $m = 1$ , then  $m - 1 = 0$  and so also  $m - 1 \leq nx$  (as  $x \geq 0$ ).

$n(y - x) > 1 \Rightarrow ny > 1 + nx \Rightarrow ny > 1 + nx \geq m \Rightarrow y > m/n$  So  $r = m/n$  works.

Now assume  $x < 0$ . If  $y > 0$ , then  $r = 0$  works.

If  $y \leq 0$ , then  $0 \leq -y < -x$ . Find  $q \in \mathbb{Q}$  s.t.  $-y < q < -x$ .  $r = -q$  works.




## Corollary

$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$

## Corollary


$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$





## Corollary


$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$



**Proof:**  $A := \{1/n : n \in \mathbb{N}\}$  is nonempty.

## Corollary

$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$




**Proof:**  $A := \{1/n : n \in \mathbb{N}\}$  is nonempty.

$1/n > 0$  for all  $n \in \mathbb{N} \Rightarrow 0$  is a lower bound

## Corollary

$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$




**Proof:**  $A := \{1/n : n \in \mathbb{N}\}$  is nonempty.

$1/n > 0$  for all  $n \in \mathbb{N} \Rightarrow 0$  is a lower bound

$\Rightarrow b := \inf A$  exists and  $b \geq 0$ .

## Corollary

$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$



**Proof:**  $A := \{1/n : n \in \mathbb{N}\}$  is nonempty.


$1/n > 0$  for all  $n \in \mathbb{N} \Rightarrow 0$  is a lower bound

$\Rightarrow b := \inf A$  exists and  $b \geq 0$ .

Let  $a > 0$  be arbitrary.

## Corollary

$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$



**Proof:**  $A := \{1/n : n \in \mathbb{N}\}$  is nonempty.

$1/n > 0$  for all  $n \in \mathbb{N} \Rightarrow 0$  is a lower bound


$\Rightarrow b := \inf A$  exists and  $b \geq 0$ .

Let  $a > 0$  be arbitrary.

Archimedean property  $\Rightarrow \exists n \in \mathbb{N}$  such that  $na > 1$

## Corollary

$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$



**Proof:**  $A := \{1/n : n \in \mathbb{N}\}$  is nonempty.

$1/n > 0$  for all  $n \in \mathbb{N} \Rightarrow 0$  is a lower bound


$\Rightarrow b := \inf A$  exists and  $b \geq 0$ .

Let  $a > 0$  be arbitrary.

Archimedean property  $\Rightarrow \exists n \in \mathbb{N}$  such that  $na > 1 \Rightarrow a > 1/n \in A$

## Corollary

$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$



**Proof:**  $A := \{1/n : n \in \mathbb{N}\}$  is nonempty.

$1/n > 0$  for all  $n \in \mathbb{N} \Rightarrow 0$  is a lower bound

$\Rightarrow b := \inf A$  exists and  $b \geq 0$ .


Let  $a > 0$  be arbitrary.

Archimedean property  $\Rightarrow \exists n \in \mathbb{N}$  such that  $na > 1 \Rightarrow a > 1/n \in A$

$\Rightarrow a$  is not a lower bound

## Corollary

$$\inf\{1/n : n \in \mathbb{N}\} = 0.$$



**Proof:**  $A := \{1/n : n \in \mathbb{N}\}$  is nonempty.

$1/n > 0$  for all  $n \in \mathbb{N} \Rightarrow 0$  is a lower bound

$\Rightarrow b := \inf A$  exists and  $b \geq 0$ .

Let  $a > 0$  be arbitrary.

Archimedean property  $\Rightarrow \exists n \in \mathbb{N}$  such that  $na > 1 \Rightarrow a > 1/n \in A$

$\Rightarrow a$  is not a lower bound  $\Rightarrow b = 0$ . □



For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

E.g., if  $A = \{1, 2, 3\}$ , then  $5 + A = \{6, 7, 8\}$  and  $3A = \{3, 6, 9\}$ .

For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

E.g., if  $A = \{1, 2, 3\}$ , then  $5 + A = \{6, 7, 8\}$  and  $3A = \{3, 6, 9\}$ .

## Proposition

Let  $A \subset \mathbb{R}$  be nonempty.

(i) If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ .

For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

E.g., if  $A = \{1, 2, 3\}$ , then  $5 + A = \{6, 7, 8\}$  and  $3A = \{3, 6, 9\}$ .

## Proposition

Let  $A \subset \mathbb{R}$  be nonempty.

- (i) If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ .
- (ii) If  $x \in \mathbb{R}$  and  $A$  is bounded below, then  $\inf(x + A) = x + \inf A$ .

For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

E.g., if  $A = \{1, 2, 3\}$ , then  $5 + A = \{6, 7, 8\}$  and  $3A = \{3, 6, 9\}$ .

## Proposition

Let  $A \subset \mathbb{R}$  be nonempty.

- (i) If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ .
- (ii) If  $x \in \mathbb{R}$  and  $A$  is bounded below, then  $\inf(x + A) = x + \inf A$ .
- (iii) If  $x > 0$  and  $A$  is bounded above, then  $\sup(xA) = x(\sup A)$ .

For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

E.g., if  $A = \{1, 2, 3\}$ , then  $5 + A = \{6, 7, 8\}$  and  $3A = \{3, 6, 9\}$ .

## Proposition

Let  $A \subset \mathbb{R}$  be nonempty.

- (i) If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ .
- (ii) If  $x \in \mathbb{R}$  and  $A$  is bounded below, then  $\inf(x + A) = x + \inf A$ .
- (iii) If  $x > 0$  and  $A$  is bounded above, then  $\sup(xA) = x(\sup A)$ .
- (iv) If  $x > 0$  and  $A$  is bounded below, then  $\inf(xA) = x(\inf A)$ .

For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

E.g., if  $A = \{1, 2, 3\}$ , then  $5 + A = \{6, 7, 8\}$  and  $3A = \{3, 6, 9\}$ .

## Proposition

Let  $A \subset \mathbb{R}$  be nonempty.

- (i) If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ .
- (ii) If  $x \in \mathbb{R}$  and  $A$  is bounded below, then  $\inf(x + A) = x + \inf A$ .
- (iii) If  $x > 0$  and  $A$  is bounded above, then  $\sup(xA) = x(\sup A)$ .
- (iv) If  $x > 0$  and  $A$  is bounded below, then  $\inf(xA) = x(\inf A)$ .
- (v) If  $x < 0$  and  $A$  is bounded below, then  $\sup(xA) = x(\inf A)$ .

For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

E.g., if  $A = \{1, 2, 3\}$ , then  $5 + A = \{6, 7, 8\}$  and  $3A = \{3, 6, 9\}$ .

## Proposition

Let  $A \subset \mathbb{R}$  be nonempty.

- (i) If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ .
- (ii) If  $x \in \mathbb{R}$  and  $A$  is bounded below, then  $\inf(x + A) = x + \inf A$ .
- (iii) If  $x > 0$  and  $A$  is bounded above, then  $\sup(xA) = x(\sup A)$ .
- (iv) If  $x > 0$  and  $A$  is bounded below, then  $\inf(xA) = x(\inf A)$ .
- (v) If  $x < 0$  and  $A$  is bounded below, then  $\sup(xA) = x(\inf A)$ .
- (vi) If  $x < 0$  and  $A$  is bounded above, then  $\inf(xA) = x(\sup A)$ .



For  $A \subset \mathbb{R}$ ,  $x \in \mathbb{R}$  define  $x + A := \{x + y \in \mathbb{R} : y \in A\}$   $xA := \{xy \in \mathbb{R} : y \in A\}$ .

E.g., if  $A = \{1, 2, 3\}$ , then  $5 + A = \{6, 7, 8\}$  and  $3A = \{3, 6, 9\}$ .

## Proposition

Let  $A \subset \mathbb{R}$  be nonempty.

- (i) If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ .
- (ii) If  $x \in \mathbb{R}$  and  $A$  is bounded below, then  $\inf(x + A) = x + \inf A$ .
- (iii) If  $x > 0$  and  $A$  is bounded above, then  $\sup(xA) = x(\sup A)$ .
- (iv) If  $x > 0$  and  $A$  is bounded below, then  $\inf(xA) = x(\inf A)$ .
- (v) If  $x < 0$  and  $A$  is bounded below, then  $\sup(xA) = x(\inf A)$ .
- (vi) If  $x < 0$  and  $A$  is bounded above, then  $\inf(xA) = x(\sup A)$ .

Note that if  $x < 0$ , then for multiplication supremum and infimum switch.

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

$$\Rightarrow x + b \text{ is an upper bound for } x + A.$$

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

$$\Rightarrow x + b \text{ is an upper bound for } x + A.$$

$$\text{If } b = \sup A, \text{ then } \sup(x + A) \leq x + b = x + \sup A.$$

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

$$\Rightarrow x + b \text{ is an upper bound for } x + A.$$

$$\text{If } b = \sup A, \text{ then } \sup(x + A) \leq x + b = x + \sup A.$$

Opposite inequality is similar:

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

$$\Rightarrow x + b \text{ is an upper bound for } x + A.$$

If  $b = \sup A$ , then  $\sup(x + A) \leq x + b = x + \sup A$ .

Opposite inequality is similar: Suppose  $c$  is an upper bound for  $x + A$ .



**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

$$\Rightarrow x + b \text{ is an upper bound for } x + A.$$

If  $b = \sup A$ , then  $\sup(x + A) \leq x + b = x + \sup A$ .

Opposite inequality is similar: Suppose  $c$  is an upper bound for  $x + A$ .

$$\Rightarrow x + y \leq c \text{ for all } y \in A$$

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

$$\Rightarrow x + b \text{ is an upper bound for } x + A.$$

If  $b = \sup A$ , then  $\sup(x + A) \leq x + b = x + \sup A$ .

Opposite inequality is similar: Suppose  $c$  is an upper bound for  $x + A$ .

$$\Rightarrow x + y \leq c \text{ for all } y \in A \quad \Rightarrow \quad y \leq c - x \text{ for all } y \in A$$

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

$$\Rightarrow x + b \text{ is an upper bound for } x + A.$$

If  $b = \sup A$ , then  $\sup(x + A) \leq x + b = x + \sup A$ .

Opposite inequality is similar: Suppose  $c$  is an upper bound for  $x + A$ .

$$\Rightarrow x + y \leq c \text{ for all } y \in A \quad \Rightarrow \quad y \leq c - x \text{ for all } y \in A$$

$$\Rightarrow c - x \text{ is an upper bound for } A.$$

**Proof:** Let's prove (i):

"If  $x \in \mathbb{R}$  and  $A$  is bounded above, then  $\sup(x + A) = x + \sup A$ ",  
rest are exercises.

Suppose  $b$  is an upper bound for  $A$ .

$$\Rightarrow y \leq b \text{ for all } y \in A \quad \Rightarrow \quad x + y \leq x + b \text{ for all } y \in A$$

$$\Rightarrow x + b \text{ is an upper bound for } x + A.$$

$$\text{If } b = \sup A, \text{ then } \sup(x + A) \leq x + b = x + \sup A.$$

Opposite inequality is similar: Suppose  $c$  is an upper bound for  $x + A$ .

$$\Rightarrow x + y \leq c \text{ for all } y \in A \quad \Rightarrow \quad y \leq c - x \text{ for all } y \in A$$

$$\Rightarrow c - x \text{ is an upper bound for } A.$$

$$\text{If } c = \sup(x + A), \text{ then } \sup A \leq c - x = \sup(x + A) - x.$$



## Proposition

*Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ .*

## Proposition

*Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ .  
Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .*

## Proposition

*Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ . Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .*

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

## Proposition

*Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ . Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .*

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

$\Rightarrow x \leq \inf B$  for all  $x \in A$



## Proposition

*Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ . Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .*

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

$\Rightarrow x \leq \inf B$  for all  $x \in A$

$\Rightarrow \inf B$  is an upper bound for  $A$

## Proposition

Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ .  
Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

$\Rightarrow x \leq \inf B$  for all  $x \in A$

$\Rightarrow \inf B$  is an upper bound for  $A$

$\Rightarrow \sup A \leq \inf B$ .



## Proposition

*Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ . Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .*

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

$\Rightarrow x \leq \inf B$  for all  $x \in A$

$\Rightarrow \inf B$  is an upper bound for  $A$

$\Rightarrow \sup A \leq \inf B$ . □

Care must be taken with suprema and infima and strict inequalities.

## Proposition

Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ .  
Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

$\Rightarrow x \leq \inf B$  for all  $x \in A$

$\Rightarrow \inf B$  is an upper bound for  $A$

$\Rightarrow \sup A \leq \inf B$ . □

Care must be taken with suprema and infima and strict inequalities.

“ $x < y$  for all  $x \in A$  and  $y \in B$ ” still only implies  $\sup A \leq \inf B$  (nonstrict).

## Proposition

Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ .  
Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

$\Rightarrow x \leq \inf B$  for all  $x \in A$

$\Rightarrow \inf B$  is an upper bound for  $A$

$\Rightarrow \sup A \leq \inf B$ . □

Care must be taken with suprema and infima and strict inequalities.

“ $x < y$  for all  $x \in A$  and  $y \in B$ ” still only implies  $\sup A \leq \inf B$  (nonstrict).

E.g.,  $A := \{0\}$  and  $B := \{1/n : n \in \mathbb{N}\}$

## Proposition

Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ .  
Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

$\Rightarrow x \leq \inf B$  for all  $x \in A$

$\Rightarrow \inf B$  is an upper bound for  $A$

$\Rightarrow \sup A \leq \inf B$ . □

Care must be taken with suprema and infima and strict inequalities.

“ $x < y$  for all  $x \in A$  and  $y \in B$ ” still only implies  $\sup A \leq \inf B$  (nonstrict).

E.g.,  $A := \{0\}$  and  $B := \{1/n : n \in \mathbb{N}\} \Rightarrow 0 < 1/n$  for all  $n \in \mathbb{N}$ .

## Proposition

Let  $A, B \subset \mathbb{R}$  be nonempty sets such that  $x \leq y$  whenever  $x \in A$  and  $y \in B$ .  
Then  $A$  is bounded above,  $B$  is bounded below, and  $\sup A \leq \inf B$ .

**Proof:** Any  $x \in A$  is a lower bound for  $B$ .

$\Rightarrow x \leq \inf B$  for all  $x \in A$

$\Rightarrow \inf B$  is an upper bound for  $A$

$\Rightarrow \sup A \leq \inf B$ . □

Care must be taken with suprema and infima and strict inequalities.

“ $x < y$  for all  $x \in A$  and  $y \in B$ ” still only implies  $\sup A \leq \inf B$  (nonstrict).

E.g.,  $A := \{0\}$  and  $B := \{1/n : n \in \mathbb{N}\} \Rightarrow 0 < 1/n$  for all  $n \in \mathbb{N}$ .

However,  $\sup A = 0$  and  $\inf B = 0$ .

## Proposition

*If  $S \subset \mathbb{R}$  is nonempty and bounded above, then for every  $\epsilon > 0$  there exists an  $x \in S$  such that  $(\sup S) - \epsilon < x \leq \sup S$ .*



## Proposition

*If  $S \subset \mathbb{R}$  is nonempty and bounded above, then for every  $\epsilon > 0$  there exists an  $x \in S$  such that  $(\sup S) - \epsilon < x \leq \sup S$ .*

Proof is an exercise.

## Definition

Let  $A \subset \mathbb{R}$  be a set.

## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .

## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .
- (ii) If  $A$  is not bounded above, then  $\sup A := \infty$ .

## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .
- (ii) If  $A$  is not bounded above, then  $\sup A := \infty$ .
- (iii) If  $A$  is empty, then  $\inf A := \infty$ .

## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .
- (ii) If  $A$  is not bounded above, then  $\sup A := \infty$ .
- (iii) If  $A$  is empty, then  $\inf A := \infty$ .
- (iv) If  $A$  is not bounded below, then  $\inf A := -\infty$ .

## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .
- (ii) If  $A$  is not bounded above, then  $\sup A := \infty$ .
- (iii) If  $A$  is empty, then  $\inf A := \infty$ .
- (iv) If  $A$  is not bounded below, then  $\inf A := -\infty$ .

**Remark:**  $\infty$  and  $-\infty$  are sometimes treated as numbers.

## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .
- (ii) If  $A$  is not bounded above, then  $\sup A := \infty$ .
- (iii) If  $A$  is empty, then  $\inf A := \infty$ .
- (iv) If  $A$  is not bounded below, then  $\inf A := -\infty$ .

**Remark:**  $\infty$  and  $-\infty$  are sometimes treated as numbers.

$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$  (extended real numbers) is an ordered set  
( $-\infty < \infty$  and  $-\infty < x$  and  $x < \infty$  for all  $x \in \mathbb{R}$ ).



## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .
- (ii) If  $A$  is not bounded above, then  $\sup A := \infty$ .
- (iii) If  $A$  is empty, then  $\inf A := \infty$ .
- (iv) If  $A$  is not bounded below, then  $\inf A := -\infty$ .

**Remark:**  $\infty$  and  $-\infty$  are sometimes treated as numbers.

$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$  (extended real numbers) is an ordered set  
( $-\infty < \infty$  and  $-\infty < x$  and  $x < \infty$  for all  $x \in \mathbb{R}$ ).

Some (but not all) arithmetic can be done in the obvious way,

## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .
- (ii) If  $A$  is not bounded above, then  $\sup A := \infty$ .
- (iii) If  $A$  is empty, then  $\inf A := \infty$ .
- (iv) If  $A$  is not bounded below, then  $\inf A := -\infty$ .

**Remark:**  $\infty$  and  $-\infty$  are sometimes treated as numbers.

$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$  (extended real numbers) is an ordered set  
( $-\infty < \infty$  and  $-\infty < x$  and  $x < \infty$  for all  $x \in \mathbb{R}$ ).

Some (but not all) arithmetic can be done in the obvious way,

leave  $\infty - \infty$ ,  $0 \cdot (\pm\infty)$ , and  $\frac{\pm\infty}{\pm\infty}$  undefined.

## Definition

Let  $A \subset \mathbb{R}$  be a set.

- (i) If  $A$  is empty, then  $\sup A := -\infty$ .
- (ii) If  $A$  is not bounded above, then  $\sup A := \infty$ .
- (iii) If  $A$  is empty, then  $\inf A := \infty$ .
- (iv) If  $A$  is not bounded below, then  $\inf A := -\infty$ .

**Remark:**  $\infty$  and  $-\infty$  are sometimes treated as numbers.

$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$  (extended real numbers) is an ordered set  
( $-\infty < \infty$  and  $-\infty < x$  and  $x < \infty$  for all  $x \in \mathbb{R}$ ).

Some (but not all) arithmetic can be done in the obvious way,  
leave  $\infty - \infty$ ,  $0 \cdot (\pm\infty)$ , and  $\frac{\pm\infty}{\pm\infty}$  undefined.

We avoid using this arithmetic;  $\mathbb{R}^*$  is not a field!

If  $A \neq \emptyset$  is finite, then  $\inf A \in A$  and  $\sup A \in A$ .

If  $A \neq \emptyset$  is finite, then  $\inf A \in A$  and  $\sup A \in A$ .

We then usually write

$\min A := \inf A$       (minimum)

$\max A := \sup A$       (maximum)

If  $A \neq \emptyset$  is finite, then  $\inf A \in A$  and  $\sup A \in A$ .

We then usually write

$\min A := \inf A$  (minimum)

$\max A := \sup A$  (maximum)

E.g.,  $\max\{1, 2.4, \pi, 100\} = 100$        $\min\{1, 2.4, \pi, 100\} = 1$ .

Some exercises with useful conclusions.

Some exercises with useful conclusions.

**Exercise:** Prove that if  $t \geq 0$  ( $t \in \mathbb{R}$ ), then there exists an  $n \in \mathbb{N}$  such that  $n - 1 \leq t < n$ .



Some exercises with useful conclusions.

**Exercise:** Prove that if  $t \geq 0$  ( $t \in \mathbb{R}$ ), then there exists an  $n \in \mathbb{N}$  such that  $n - 1 \leq t < n$ .

**Exercise:** Let  $x, y \in \mathbb{R}$ . Suppose  $x^2 + y^2 = 0$ . Prove that  $x = 0$  and  $y = 0$ .

Some exercises with useful conclusions.

**Exercise:** Prove that if  $t \geq 0$  ( $t \in \mathbb{R}$ ), then there exists an  $n \in \mathbb{N}$  such that  $n - 1 \leq t < n$ .

**Exercise:** Let  $x, y \in \mathbb{R}$ . Suppose  $x^2 + y^2 = 0$ . Prove that  $x = 0$  and  $y = 0$ .

**Exercise:** Let  $A$  and  $B$  be two nonempty bounded sets of real numbers. Let  $C := \{a + b : a \in A, b \in B\}$ . Show that  $C$  is a bounded set and that

$$\sup C = \sup A + \sup B \quad \text{and} \quad \inf C = \inf A + \inf B.$$

Some exercises with useful conclusions.

**Exercise:** Prove that if  $t \geq 0$  ( $t \in \mathbb{R}$ ), then there exists an  $n \in \mathbb{N}$  such that  $n - 1 \leq t < n$ .

**Exercise:** Let  $x, y \in \mathbb{R}$ . Suppose  $x^2 + y^2 = 0$ . Prove that  $x = 0$  and  $y = 0$ .

**Exercise:** Let  $A$  and  $B$  be two nonempty bounded sets of real numbers. Let  $C := \{a + b : a \in A, b \in B\}$ . Show that  $C$  is a bounded set and that

$$\sup C = \sup A + \sup B \quad \text{and} \quad \inf C = \inf A + \inf B.$$

**Exercise:** Prove the so-called *Bernoulli's inequality*: If  $1 + x > 0$ , then for all  $n \in \mathbb{N}$ , we have  $(1 + x)^n \geq 1 + nx$ .