

BA: 3.2

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

Let $S \subset \mathbb{R}$, $c \in S$. We say $f: S \rightarrow \mathbb{R}$ is *continuous at c* if for every $\epsilon > 0$ there is a $\delta > 0$ such that whenever $x \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

Definition


Let $S \subset \mathbb{R}$, $c \in S$. We say $f: S \rightarrow \mathbb{R}$ is *continuous at c* if for every $\epsilon > 0$ there is a $\delta > 0$ such that whenever $x \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

When $f: S \rightarrow \mathbb{R}$ is continuous at all $c \in S$, we say f is a *continuous function*.

Definition

Let $S \subset \mathbb{R}$, $c \in S$. We say $f: S \rightarrow \mathbb{R}$ is *continuous at c* if for every $\epsilon > 0$ there is a $\delta > 0$ such that whenever $x \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.


When $f: S \rightarrow \mathbb{R}$ is continuous at all $c \in S$, we say f is a *continuous function*.



Definition

Let $S \subset \mathbb{R}$, $c \in S$. We say $f: S \rightarrow \mathbb{R}$ is *continuous at c* if for every $\epsilon > 0$ there is a $\delta > 0$ such that whenever $x \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

When $f: S \rightarrow \mathbb{R}$ is continuous at all $c \in S$, we say f is a *continuous function*.




Note: δ depends on both ϵ and c ; no need to pick the same δ for all $c \in S$.

Definition

Let $S \subset \mathbb{R}$, $c \in S$. We say $f: S \rightarrow \mathbb{R}$ is *continuous at c* if for every $\epsilon > 0$ there is a $\delta > 0$ such that whenever $x \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

When $f: S \rightarrow \mathbb{R}$ is continuous at all $c \in S$, we say f is a *continuous function*.




Note: δ depends on both ϵ and c ; no need to pick the same δ for all $c \in S$.

If f is continuous for all $c \in A$, we say f is *continuous on $A \subset S$* .

Definition

Let $S \subset \mathbb{R}$, $c \in S$. We say $f: S \rightarrow \mathbb{R}$ is *continuous at c* if for every $\epsilon > 0$ there is a $\delta > 0$ such that whenever $x \in S$ and $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

When $f: S \rightarrow \mathbb{R}$ is continuous at all $c \in S$, we say f is a *continuous function*.



Note: δ depends on both ϵ and c ; no need to pick the same δ for all $c \in S$.

If f is continuous for all $c \in A$, we say f is *continuous on $A \subset S$* .

Remark: If f is continuous on A , then $f|_A$ is continuous (exercise), but the converse does not hold (we'll give an example shortly).

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $S \cap (c - \delta, c + \delta) = \{c\}$.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $S \cap (c - \delta, c + \delta) = \{c\}$.

If $x \in S$ and $|x - c| < \delta$, then $x = c$.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $S \cap (c - \delta, c + \delta) = \{c\}$.

If $x \in S$ and $|x - c| < \delta$, then $x = c$. $\Rightarrow |f(x) - f(c)| = |f(c) - f(c)| = 0 < \epsilon$.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $S \cap (c - \delta, c + \delta) = \{c\}$.

If $x \in S$ and $|x - c| < \delta$, then $x = c$. $\Rightarrow |f(x) - f(c)| = |f(c) - f(c)| = 0 < \epsilon$.

(ii) Suppose c is a cluster point of S .

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $S \cap (c - \delta, c + \delta) = \{c\}$.

If $x \in S$ and $|x - c| < \delta$, then $x = c$. $\Rightarrow |f(x) - f(c)| = |f(c) - f(c)| = 0 < \epsilon$.

(ii) Suppose c is a cluster point of S .

First suppose $\lim_{x \rightarrow c} f(x) = f(c)$.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $S \cap (c - \delta, c + \delta) = \{c\}$.

If $x \in S$ and $|x - c| < \delta$, then $x = c$. $\Rightarrow |f(x) - f(c)| = |f(c) - f(c)| = 0 < \epsilon$.

(ii) Suppose c is a cluster point of S .

First suppose $\lim_{x \rightarrow c} f(x) = f(c)$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $x \in S \setminus \{c\}$ and $|x - c| < \delta$, then $|f(x) - f(c)| < \epsilon$.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $S \cap (c - \delta, c + \delta) = \{c\}$.

If $x \in S$ and $|x - c| < \delta$, then $x = c$. $\Rightarrow |f(x) - f(c)| = |f(c) - f(c)| = 0 < \epsilon$.

(ii) Suppose c is a cluster point of S .

First suppose $\lim_{x \rightarrow c} f(x) = f(c)$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $x \in S \setminus \{c\}$ and $|x - c| < \delta$, then $|f(x) - f(c)| < \epsilon$.

Also $|f(c) - f(c)| = 0 < \epsilon$.

Proposition

Consider $f: S \rightarrow \mathbb{R}$ where $S \subset \mathbb{R}$ and let $c \in S$. Then:

- (i) If c is not a cluster point of S , then f is continuous at c .
- (ii) If c is a cluster point of S , then f is continuous at c if and only if the limit of $f(x)$ as $x \rightarrow c$ exists and

$$\lim_{x \rightarrow c} f(x) = f(c).$$

- (iii) The function f is continuous at c if and only if for every sequence $\{x_n\}_{n=1}^{\infty}$ where $x_n \in S$ and $\lim_{n \rightarrow \infty} x_n = c$, the sequence $\{f(x_n)\}_{n=1}^{\infty}$ converges to $f(c)$.

Proof: (i) Suppose c is not a cluster point of S . Let $\epsilon > 0$ be given.

$\exists \delta > 0$ such that $S \cap (c - \delta, c + \delta) = \{c\}$.

If $x \in S$ and $|x - c| < \delta$, then $x = c$. $\Rightarrow |f(x) - f(c)| = |f(c) - f(c)| = 0 < \epsilon$.

(ii) Suppose c is a cluster point of S .

First suppose $\lim_{x \rightarrow c} f(x) = f(c)$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. if $x \in S \setminus \{c\}$ and $|x - c| < \delta$, then $|f(x) - f(c)| < \epsilon$.

Also $|f(c) - f(c)| = 0 < \epsilon$. $\Rightarrow f$ is continuous at c .

Now suppose f is continuous at c .

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

$\exists M \in \mathbb{N}$ such that $\forall n \geq M$, we have $|x_n - c| < \delta$.

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

$\exists M \in \mathbb{N}$ such that $\forall n \geq M$, we have $|x_n - c| < \delta$.

$$\Rightarrow \text{if } n \geq M, \text{ then } |f(x_n) - f(c)| < \epsilon.$$

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

$\exists M \in \mathbb{N}$ such that $\forall n \geq M$, we have $|x_n - c| < \delta$.

$$\Rightarrow \text{if } n \geq M, \text{ then } |f(x_n) - f(c)| < \epsilon. \quad \Rightarrow \quad \{f(x_n)\}_{n=1}^{\infty} \text{ converges to } f(c).$$

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

$\exists M \in \mathbb{N}$ such that $\forall n \geq M$, we have $|x_n - c| < \delta$.

$$\Rightarrow \text{if } n \geq M, \text{ then } |f(x_n) - f(c)| < \epsilon. \quad \Rightarrow \quad \{f(x_n)\}_{n=1}^{\infty} \text{ converges to } f(c).$$

Suppose f is not continuous at c .

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

$\exists M \in \mathbb{N}$ such that $\forall n \geq M$, we have $|x_n - c| < \delta$.

$$\Rightarrow \text{if } n \geq M, \text{ then } |f(x_n) - f(c)| < \epsilon. \quad \Rightarrow \quad \{f(x_n)\}_{n=1}^{\infty} \text{ converges to } f(c).$$

Suppose f is not continuous at c .

$\exists \epsilon > 0$, s.t. $\forall n \in \mathbb{N}$, $\exists x_n \in S$ where $|x_n - c| < 1/n$ and $|f(x_n) - f(c)| \geq \epsilon$.

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

$\exists M \in \mathbb{N}$ such that $\forall n \geq M$, we have $|x_n - c| < \delta$.

$$\Rightarrow \text{if } n \geq M, \text{ then } |f(x_n) - f(c)| < \epsilon. \quad \Rightarrow \quad \{f(x_n)\}_{n=1}^{\infty} \text{ converges to } f(c).$$

Suppose f is not continuous at c .

$\exists \epsilon > 0$, s.t. $\forall n \in \mathbb{N}$, $\exists x_n \in S$ where $|x_n - c| < 1/n$ and $|f(x_n) - f(c)| \geq \epsilon$.

$$\lim_{n \rightarrow \infty} x_n = c, \quad \text{but as } |f(x_n) - f(c)| \geq \epsilon \quad \forall n \in \mathbb{N},$$

Now suppose f is continuous at c .

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

That's still true if $x \in S \setminus \{c\} \subset S$.

$$\Rightarrow \lim_{x \rightarrow c} f(x) = f(c).$$

(iii) First suppose f is continuous at c .

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in S and $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, $\exists \delta > 0$ s.t. $\forall x \in S$ where $|x - c| < \delta$, we have $|f(x) - f(c)| < \epsilon$.

$\exists M \in \mathbb{N}$ such that $\forall n \geq M$, we have $|x_n - c| < \delta$.

$$\Rightarrow \text{if } n \geq M, \text{ then } |f(x_n) - f(c)| < \epsilon. \quad \Rightarrow \quad \{f(x_n)\}_{n=1}^{\infty} \text{ converges to } f(c).$$

Suppose f is not continuous at c .

$\exists \epsilon > 0$, s.t. $\forall n \in \mathbb{N}$, $\exists x_n \in S$ where $|x_n - c| < 1/n$ and $|f(x_n) - f(c)| \geq \epsilon$.

$\lim_{n \rightarrow \infty} x_n = c$, but as $|f(x_n) - f(c)| \geq \epsilon \forall n \in \mathbb{N}$,

$$\Rightarrow \{f(x_n)\}_{n=1}^{\infty} \text{ does not converge to } f(c).$$



Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Proof: Fix $c \in (0, \infty)$.

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Proof: Fix $c \in (0, \infty)$.

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Proof: Fix $c \in (0, \infty)$.

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then

$$f(c) = \frac{1}{c}$$

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Proof: Fix $c \in (0, \infty)$.

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then

$$f(c) = \frac{1}{c} = \frac{1}{\lim_{n \rightarrow \infty} x_n}$$

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Proof: Fix $c \in (0, \infty)$.

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then

$$f(c) = \frac{1}{c} = \frac{1}{\lim_{n \rightarrow \infty} x_n} = \lim_{n \rightarrow \infty} \frac{1}{x_n}$$

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Proof: Fix $c \in (0, \infty)$.

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then

$$f(c) = \frac{1}{c} = \frac{1}{\lim_{n \rightarrow \infty} x_n} = \lim_{n \rightarrow \infty} \frac{1}{x_n} = \lim_{n \rightarrow \infty} f(x_n).$$

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Proof: Fix $c \in (0, \infty)$.

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then

$$f(c) = \frac{1}{c} = \frac{1}{\lim_{n \rightarrow \infty} x_n} = \lim_{n \rightarrow \infty} \frac{1}{x_n} = \lim_{n \rightarrow \infty} f(x_n).$$

$\Rightarrow f$ is continuous at c .

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ defined by $f(x) := 1/x$ is continuous.

Proof: Fix $c \in (0, \infty)$.

Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in $(0, \infty)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then

$$f(c) = \frac{1}{c} = \frac{1}{\lim_{n \rightarrow \infty} x_n} = \lim_{n \rightarrow \infty} \frac{1}{x_n} = \lim_{n \rightarrow \infty} f(x_n).$$

$\Rightarrow f$ is continuous at c .

As f is continuous at all $c \in (0, \infty)$, f is continuous.



Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proof: Fix $c \in \mathbb{R}$.

Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proof: Fix $c \in \mathbb{R}$. Let $\{x_n\}_{n=1}^{\infty}$ be such that $\lim_{n \rightarrow \infty} x_n = c$.

Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proof: Fix $c \in \mathbb{R}$. Let $\{x_n\}_{n=1}^{\infty}$ be such that $\lim_{n \rightarrow \infty} x_n = c$. Then

$$f(c) = a_d c^d + a_{d-1} c^{d-1} + \cdots + a_1 c + a_0$$

Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proof: Fix $c \in \mathbb{R}$. Let $\{x_n\}_{n=1}^{\infty}$ be such that $\lim_{n \rightarrow \infty} x_n = c$. Then

$$\begin{aligned} f(c) &= a_d c^d + a_{d-1} c^{d-1} + \cdots + a_1 c + a_0 \\ &= a_d \left(\lim_{n \rightarrow \infty} x_n \right)^d + a_{d-1} \left(\lim_{n \rightarrow \infty} x_n \right)^{d-1} + \cdots + a_1 \left(\lim_{n \rightarrow \infty} x_n \right) + a_0 \end{aligned}$$

Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proof: Fix $c \in \mathbb{R}$. Let $\{x_n\}_{n=1}^{\infty}$ be such that $\lim_{n \rightarrow \infty} x_n = c$. Then

$$\begin{aligned} f(c) &= a_d c^d + a_{d-1} c^{d-1} + \cdots + a_1 c + a_0 \\ &= a_d \left(\lim_{n \rightarrow \infty} x_n \right)^d + a_{d-1} \left(\lim_{n \rightarrow \infty} x_n \right)^{d-1} + \cdots + a_1 \left(\lim_{n \rightarrow \infty} x_n \right) + a_0 \\ &= \lim_{n \rightarrow \infty} \left(a_d x_n^d + a_{d-1} x_n^{d-1} + \cdots + a_1 x_n + a_0 \right) \end{aligned}$$

Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proof: Fix $c \in \mathbb{R}$. Let $\{x_n\}_{n=1}^\infty$ be such that $\lim_{n \rightarrow \infty} x_n = c$. Then

$$\begin{aligned} f(c) &= a_d c^d + a_{d-1} c^{d-1} + \cdots + a_1 c + a_0 \\ &= a_d \left(\lim_{n \rightarrow \infty} x_n \right)^d + a_{d-1} \left(\lim_{n \rightarrow \infty} x_n \right)^{d-1} + \cdots + a_1 \left(\lim_{n \rightarrow \infty} x_n \right) + a_0 \\ &= \lim_{n \rightarrow \infty} \left(a_d x_n^d + a_{d-1} x_n^{d-1} + \cdots + a_1 x_n + a_0 \right) = \lim_{n \rightarrow \infty} f(x_n). \end{aligned}$$

Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proof: Fix $c \in \mathbb{R}$. Let $\{x_n\}_{n=1}^\infty$ be such that $\lim_{n \rightarrow \infty} x_n = c$. Then

$$\begin{aligned} f(c) &= a_d c^d + a_{d-1} c^{d-1} + \cdots + a_1 c + a_0 \\ &= a_d \left(\lim_{n \rightarrow \infty} x_n \right)^d + a_{d-1} \left(\lim_{n \rightarrow \infty} x_n \right)^{d-1} + \cdots + a_1 \left(\lim_{n \rightarrow \infty} x_n \right) + a_0 \\ &= \lim_{n \rightarrow \infty} \left(a_d x_n^d + a_{d-1} x_n^{d-1} + \cdots + a_1 x_n + a_0 \right) = \lim_{n \rightarrow \infty} f(x_n). \end{aligned}$$

$\Rightarrow f$ is continuous at c

Proposition

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a polynomial. That is

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0,$$

for some constants a_0, a_1, \dots, a_d . Then f is continuous.

Proof: Fix $c \in \mathbb{R}$. Let $\{x_n\}_{n=1}^{\infty}$ be such that $\lim_{n \rightarrow \infty} x_n = c$. Then

$$\begin{aligned} f(c) &= a_d c^d + a_{d-1} c^{d-1} + \cdots + a_1 c + a_0 \\ &= a_d \left(\lim_{n \rightarrow \infty} x_n \right)^d + a_{d-1} \left(\lim_{n \rightarrow \infty} x_n \right)^{d-1} + \cdots + a_1 \left(\lim_{n \rightarrow \infty} x_n \right) + a_0 \\ &= \lim_{n \rightarrow \infty} \left(a_d x_n^d + a_{d-1} x_n^{d-1} + \cdots + a_1 x_n + a_0 \right) = \lim_{n \rightarrow \infty} f(x_n). \end{aligned}$$

$\Rightarrow f$ is continuous at $c \quad \Rightarrow f$ is continuous (c was arbitrary).



Proposition

Let $f: S \rightarrow \mathbb{R}$ and $g: S \rightarrow \mathbb{R}$ be functions continuous at $c \in S$.

Proposition

Let $f: S \rightarrow \mathbb{R}$ and $g: S \rightarrow \mathbb{R}$ be functions continuous at $c \in S$.

- (i) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) + g(x)$ is continuous at c .

Proposition

Let $f: S \rightarrow \mathbb{R}$ and $g: S \rightarrow \mathbb{R}$ be functions continuous at $c \in S$.

- (i) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) + g(x)$ is continuous at c .
- (ii) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) - g(x)$ is continuous at c .

Proposition

Let $f: S \rightarrow \mathbb{R}$ and $g: S \rightarrow \mathbb{R}$ be functions continuous at $c \in S$.

- (i) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) + g(x)$ is continuous at c .
- (ii) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) - g(x)$ is continuous at c .
- (iii) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x)g(x)$ is continuous at c .

Proposition

Let $f: S \rightarrow \mathbb{R}$ and $g: S \rightarrow \mathbb{R}$ be functions continuous at $c \in S$.

- (i) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) + g(x)$ is continuous at c .
- (ii) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) - g(x)$ is continuous at c .
- (iii) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x)g(x)$ is continuous at c .
- (iv) If $g(x) \neq 0$ for all $x \in S$, the function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := \frac{f(x)}{g(x)}$ is continuous at c .

Proposition

Let $f: S \rightarrow \mathbb{R}$ and $g: S \rightarrow \mathbb{R}$ be functions continuous at $c \in S$.

- (i) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) + g(x)$ is continuous at c .
- (ii) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x) - g(x)$ is continuous at c .
- (iii) The function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := f(x)g(x)$ is continuous at c .
- (iv) If $g(x) \neq 0$ for all $x \in S$, the function $h: S \rightarrow \mathbb{R}$ defined by $h(x) := \frac{f(x)}{g(x)}$ is continuous at c .

Proof: Exercise.

Example: $\sin(x)$ and $\cos(x)$ are continuous.

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$|\sin(x) - \sin(c)|$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$|\sin(x) - \sin(c)| = \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right|$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$\begin{aligned} |\sin(x) - \sin(c)| &= \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right| \\ &= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right| \end{aligned}$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$\begin{aligned} |\sin(x) - \sin(c)| &= \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right| \\ &= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right| \\ &\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \end{aligned}$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$\begin{aligned} |\sin(x) - \sin(c)| &= \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right| \\ &= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right| \\ &\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c| \end{aligned}$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$\begin{aligned} |\sin(x) - \sin(c)| &= \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right| \\ &= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right| \\ &\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c| \end{aligned}$$

$$|\cos(x) - \cos(c)|$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$\begin{aligned} |\sin(x) - \sin(c)| &= \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right| \\ &= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right| \\ &\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c| \\ |\cos(x) - \cos(c)| &= \left| -2 \sin\left(\frac{x-c}{2}\right) \sin\left(\frac{x+c}{2}\right) \right| \end{aligned}$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$\begin{aligned} |\sin(x) - \sin(c)| &= \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right| \\ &= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right| \\ &\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c| \\ |\cos(x) - \cos(c)| &= \left| -2 \sin\left(\frac{x-c}{2}\right) \sin\left(\frac{x+c}{2}\right) \right| \\ &= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \sin\left(\frac{x+c}{2}\right) \right| \end{aligned}$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$|\sin(x) - \sin(c)| = \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right|$$

$$= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right|$$

$$\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c|$$

$$|\cos(x) - \cos(c)| = \left| -2 \sin\left(\frac{x-c}{2}\right) \sin\left(\frac{x+c}{2}\right) \right|$$

$$= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \sin\left(\frac{x+c}{2}\right) \right|$$

$$\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right|$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$|\sin(x) - \sin(c)| = \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right|$$

$$= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right|$$

$$\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c|$$

$$|\cos(x) - \cos(c)| = \left| -2 \sin\left(\frac{x-c}{2}\right) \sin\left(\frac{x+c}{2}\right) \right|$$

$$= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \sin\left(\frac{x+c}{2}\right) \right|$$

$$\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c|$$

Example: $\sin(x)$ and $\cos(x)$ are continuous.

$$|\sin(x) - \sin(c)| = \left| 2 \sin\left(\frac{x-c}{2}\right) \cos\left(\frac{x+c}{2}\right) \right|$$

$$= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \cos\left(\frac{x+c}{2}\right) \right|$$

$$\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c|$$

$$|\cos(x) - \cos(c)| = \left| -2 \sin\left(\frac{x-c}{2}\right) \sin\left(\frac{x+c}{2}\right) \right|$$

$$= 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \left| \sin\left(\frac{x+c}{2}\right) \right|$$

$$\leq 2 \left| \sin\left(\frac{x-c}{2}\right) \right| \leq 2 \left| \frac{x-c}{2} \right| = |x-c|$$

Details left to student.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

g continuous at $c \quad \Rightarrow \quad \{g(x_n)\}_{n=1}^{\infty}$ converges to $g(c)$.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

g continuous at $c \quad \Rightarrow \quad \{g(x_n)\}_{n=1}^{\infty}$ converges to $g(c)$.

f continuous at $g(c) \quad \Rightarrow \quad \{f(g(x_n))\}_{n=1}^{\infty}$ converges to $f(g(c))$.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

g continuous at $c \Rightarrow \{g(x_n)\}_{n=1}^{\infty}$ converges to $g(c)$.

f continuous at $g(c) \Rightarrow \{f(g(x_n))\}_{n=1}^{\infty}$ converges to $f(g(c))$.

$\Rightarrow f \circ g$ is continuous at c .



Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

g continuous at $c \Rightarrow \{g(x_n)\}_{n=1}^{\infty}$ converges to $g(c)$.

f continuous at $g(c) \Rightarrow \{f(g(x_n))\}_{n=1}^{\infty}$ converges to $f(g(c))$.

$\Rightarrow f \circ g$ is continuous at c . □

Example: $(\sin(1/x))^2$ is a continuous function on $(0, \infty)$.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

g continuous at $c \Rightarrow \{g(x_n)\}_{n=1}^{\infty}$ converges to $g(c)$.

f continuous at $g(c) \Rightarrow \{f(g(x_n))\}_{n=1}^{\infty}$ converges to $f(g(c))$.

$\Rightarrow f \circ g$ is continuous at c . □

Example: $(\sin(1/x))^2$ is a continuous function on $(0, \infty)$.

Proof: $1/x$ is continuous on $(0, \infty)$ and $\sin(x)$ is continuous on $(0, \infty)$.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

g continuous at $c \Rightarrow \{g(x_n)\}_{n=1}^{\infty}$ converges to $g(c)$.

f continuous at $g(c) \Rightarrow \{f(g(x_n))\}_{n=1}^{\infty}$ converges to $f(g(c))$.

$\Rightarrow f \circ g$ is continuous at c . □

Example: $(\sin(1/x))^2$ is a continuous function on $(0, \infty)$.

Proof: $1/x$ is continuous on $(0, \infty)$ and $\sin(x)$ is continuous on $(0, \infty)$.

\Rightarrow The composition $\sin(1/x)$ is continuous on $(0, \infty)$.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

g continuous at $c \Rightarrow \{g(x_n)\}_{n=1}^{\infty}$ converges to $g(c)$.

f continuous at $g(c) \Rightarrow \{f(g(x_n))\}_{n=1}^{\infty}$ converges to $f(g(c))$.

$\Rightarrow f \circ g$ is continuous at c . □

Example: $(\sin(1/x))^2$ is a continuous function on $(0, \infty)$.

Proof: $1/x$ is continuous on $(0, \infty)$ and $\sin(x)$ is continuous on $(0, \infty)$.

\Rightarrow The composition $\sin(1/x)$ is continuous on $(0, \infty)$.

x^2 is continuous on $[-1, 1]$.

Recall $f \circ g$ is defined by $(f \circ g)(x) := f(g(x))$.

Proposition

Let $A, B \subset \mathbb{R}$ and $f: B \rightarrow \mathbb{R}$ and $g: A \rightarrow B$ be functions. If g is continuous at $c \in A$ and f is continuous at $g(c)$, then $f \circ g: A \rightarrow \mathbb{R}$ is continuous at c .

Proof: Let $\{x_n\}_{n=1}^{\infty}$ be a sequence in A such that $\lim_{n \rightarrow \infty} x_n = c$.

g continuous at $c \Rightarrow \{g(x_n)\}_{n=1}^{\infty}$ converges to $g(c)$.

f continuous at $g(c) \Rightarrow \{f(g(x_n))\}_{n=1}^{\infty}$ converges to $f(g(c))$.

$\Rightarrow f \circ g$ is continuous at c . □

Example: $(\sin(1/x))^2$ is a continuous function on $(0, \infty)$.

Proof: $1/x$ is continuous on $(0, \infty)$ and $\sin(x)$ is continuous on $(0, \infty)$.

\Rightarrow The composition $\sin(1/x)$ is continuous on $(0, \infty)$.

x^2 is continuous on $[-1, 1]$. \Rightarrow The composition $(\sin(1/x))^2$ is continuous on $(0, \infty)$. □

If f is not continuous at c , say f is *discontinuous* at c (has a *discontinuity* at c).

If f is not continuous at c , say f is *discontinuous* at c (has a *discontinuity* at c).

Proposition

Consider $f: S \rightarrow \mathbb{R}$ and $c \in S$. Suppose \exists a sequence $\{x_n\}_{n=1}^{\infty}$ in S where $\lim_{n \rightarrow \infty} x_n = c$ such that $\{f(x_n)\}_{n=1}^{\infty}$ does not converge to $f(c)$.

If f is not continuous at c , say f is *discontinuous* at c (has a *discontinuity* at c).

Proposition

Consider $f: S \rightarrow \mathbb{R}$ and $c \in S$. Suppose \exists a sequence $\{x_n\}_{n=1}^{\infty}$ in S where $\lim_{n \rightarrow \infty} x_n = c$ such that $\{f(x_n)\}_{n=1}^{\infty}$ does not converge to $f(c)$.

Then f is discontinuous at c .

If f is not continuous at c , say f is *discontinuous* at c (has a *discontinuity* at c).

Proposition

Consider $f: S \rightarrow \mathbb{R}$ and $c \in S$. Suppose \exists a sequence $\{x_n\}_{n=1}^{\infty}$ in S where $\lim_{n \rightarrow \infty} x_n = c$ such that $\{f(x_n)\}_{n=1}^{\infty}$ does not converge to $f(c)$.

Then f is discontinuous at c .

Proof: A restatement of one direction of part (iii) of the proposition above. □

Example: $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) := \begin{cases} -1 & \text{if } x < 0, \\ 1 & \text{if } x \geq 0 \end{cases}$ is not continuous at 0.

Example: $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) := \begin{cases} -1 & \text{if } x < 0, \\ 1 & \text{if } x \geq 0 \end{cases}$ is not continuous at 0.

Proof: $f(-1/n) = -1 \forall n$, so $\lim_{n \rightarrow \infty} f(-1/n) = -1$, but $f(0) = 1$.

Example: $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) := \begin{cases} -1 & \text{if } x < 0, \\ 1 & \text{if } x \geq 0 \end{cases}$ is not continuous at 0.

Proof: $f(-1/n) = -1 \forall n$, so $\lim_{n \rightarrow \infty} f(-1/n) = -1$, but $f(0) = 1$.


$\Rightarrow f$ has a discontinuity at 0.



Example: $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) := \begin{cases} -1 & \text{if } x < 0, \\ 1 & \text{if } x \geq 0 \end{cases}$ is not continuous at 0.

Proof: $f(-1/n) = -1 \forall n$, so $\lim_{n \rightarrow \infty} f(-1/n) = -1$, but $f(0) = 1$.


$\Rightarrow f$ has a discontinuity at 0.



Example: $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) := \begin{cases} -1 & \text{if } x < 0, \\ 1 & \text{if } x \geq 0 \end{cases}$ is not continuous at 0.

Proof: $f(-1/n) = -1 \forall n$, so $\lim_{n \rightarrow \infty} f(-1/n) = -1$, but $f(0) = 1$.

$\Rightarrow f$ has a discontinuity at 0.




Note also:

$$f(1/n) = 1 \text{ for all } n \in \mathbb{N} \quad \Rightarrow \quad \lim_{n \rightarrow \infty} f(1/n) = f(0) = 1.$$

Example: $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) := \begin{cases} -1 & \text{if } x < 0, \\ 1 & \text{if } x \geq 0 \end{cases}$ is not continuous at 0.

Proof: $f(-1/n) = -1 \forall n$, so $\lim_{n \rightarrow \infty} f(-1/n) = -1$, but $f(0) = 1$.

$\Rightarrow f$ has a discontinuity at 0. □



Note also:

$$f(1/n) = 1 \text{ for all } n \in \mathbb{N} \quad \Rightarrow \quad \lim_{n \rightarrow \infty} f(1/n) = f(0) = 1.$$

$$f\left(\frac{(-1)^n}{n}\right) = (-1)^n.$$

Example: Consider the *Dirichlet function*.

$$f(x) := \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

Example: Consider the *Dirichlet function*.

$$f(x) := \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

f is discontinuous at all $c \in \mathbb{R}$.

Example: Consider the *Dirichlet function*.

$$f(x) := \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

f is discontinuous at all $c \in \mathbb{R}$.

Proof: For $c \in \mathbb{Q}$, take $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Example: Consider the *Dirichlet function*.

$$f(x) := \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

f is discontinuous at all $c \in \mathbb{R}$.

Proof: For $c \in \mathbb{Q}$, take $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

$f(x_n) = 0 \Rightarrow \lim_{n \rightarrow \infty} f(x_n) = 0$, but $f(c) = 1$.

Example: Consider the *Dirichlet function*.

$$f(x) := \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

f is discontinuous at all $c \in \mathbb{R}$.

Proof: For $c \in \mathbb{Q}$, take $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

$$f(x_n) = 0 \Rightarrow \lim_{n \rightarrow \infty} f(x_n) = 0, \quad \text{but } f(c) = 1.$$

For $c \in \mathbb{R} \setminus \mathbb{Q}$, take $\{x_n\}_{n=1}^{\infty}$ in \mathbb{Q} such that $\lim_{n \rightarrow \infty} x_n = c$.

Example: Consider the *Dirichlet function*.

$$f(x) := \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

f is discontinuous at all $c \in \mathbb{R}$.

Proof: For $c \in \mathbb{Q}$, take $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

$$f(x_n) = 0 \Rightarrow \lim_{n \rightarrow \infty} f(x_n) = 0, \quad \text{but } f(c) = 1.$$

For $c \in \mathbb{R} \setminus \mathbb{Q}$, take $\{x_n\}_{n=1}^{\infty}$ in \mathbb{Q} such that $\lim_{n \rightarrow \infty} x_n = c$.

$$f(x_n) = 1 \Rightarrow \lim_{n \rightarrow \infty} f(x_n) = 1, \quad \text{but } f(c) = 0.$$



Q: Does there exist a function continuous at all irrational numbers, but discontinuous at all rational numbers?

Q: Does there exist a function continuous at all irrational numbers, but discontinuous at all rational numbers?

A: Yes.


Q: Does there exist a function continuous at all irrational numbers, but discontinuous at all rational numbers?

A: Yes.

Example: (*Thomae function* or *popcorn function*).

Define $f: (0, 1) \rightarrow \mathbb{R}$ as

$$f(x) := \begin{cases} 1/k & \text{if } x = m/k, \text{ where } m, k \in \mathbb{N} \text{ and } m \text{ and } k \text{ have no common divisors,} \\ 0 & \text{if } x \text{ is irrational.} \end{cases}$$




Q: Does there exist a function continuous at all irrational numbers, but discontinuous at all rational numbers?

A: Yes.

Example: (*Thomae function* or *popcorn function*).

Define $f: (0, 1) \rightarrow \mathbb{R}$ as

$$f(x) := \begin{cases} 1/k & \text{if } x = m/k, \text{ where } m, k \in \mathbb{N} \text{ and } m \text{ and } k \text{ have no common divisors,} \\ 0 & \text{if } x \text{ is irrational.} \end{cases}$$



Claim: f is continuous at every irrational number and discontinuous at every rational number.

Proof of claim: Suppose $c = m/k$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

If $m/k \in (0, 1)$, then $0 < m < k$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

If $m/k \in (0, 1)$, then $0 < m < k$.

Only finitely many such numbers where $k < K$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

If $m/k \in (0, 1)$, then $0 < m < k$.

Only finitely many such numbers where $k < K$.

As $\lim_{n \rightarrow \infty} x_n = c$, any $x \neq c$ appears at most finitely many times in $\{x_n\}_{n=1}^{\infty}$.

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

If $m/k \in (0, 1)$, then $0 < m < k$.

Only finitely many such numbers where $k < K$.

As $\lim_{n \rightarrow \infty} x_n = c$, any $x \neq c$ appears at most finitely many times in $\{x_n\}_{n=1}^{\infty}$.

Hence, $\exists M$ such that for $n \geq M$, all the numbers x_n that are rational have a denominator larger than or equal to K .

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

If $m/k \in (0, 1)$, then $0 < m < k$.

Only finitely many such numbers where $k < K$.

As $\lim_{n \rightarrow \infty} x_n = c$, any $x \neq c$ appears at most finitely many times in $\{x_n\}_{n=1}^{\infty}$.

Hence, $\exists M$ such that for $n \geq M$, all the numbers x_n that are rational have a denominator larger than or equal to K .

$$\text{For } n \geq M, \quad |f(x_n) - 0| = f(x_n)$$

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

If $m/k \in (0, 1)$, then $0 < m < k$.

Only finitely many such numbers where $k < K$.

As $\lim_{n \rightarrow \infty} x_n = c$, any $x \neq c$ appears at most finitely many times in $\{x_n\}_{n=1}^{\infty}$.

Hence, $\exists M$ such that for $n \geq M$, all the numbers x_n that are rational have a denominator larger than or equal to K .

$$\text{For } n \geq M, \quad |f(x_n) - 0| = f(x_n) \leq 1/K$$

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

If $m/k \in (0, 1)$, then $0 < m < k$.

Only finitely many such numbers where $k < K$.

As $\lim_{n \rightarrow \infty} x_n = c$, any $x \neq c$ appears at most finitely many times in $\{x_n\}_{n=1}^{\infty}$.

Hence, $\exists M$ such that for $n \geq M$, all the numbers x_n that are rational have a denominator larger than or equal to K .

$$\text{For } n \geq M, \quad |f(x_n) - 0| = f(x_n) \leq 1/K < \epsilon.$$

Proof of claim: Suppose $c = m/k$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $\mathbb{R} \setminus \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Then $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} 0 = 0$, but $f(c) = 1/k \neq 0$.

$\Rightarrow f$ is discontinuous at c .

Now let c be irrational, so $f(c) = 0$.

Take a sequence $\{x_n\}_{n=1}^{\infty}$ in $(0, 1)$ such that $\lim_{n \rightarrow \infty} x_n = c$.

Given $\epsilon > 0$, find $K \in \mathbb{N}$ such that $1/K < \epsilon$.

If $m/k \in (0, 1)$, then $0 < m < k$.

Only finitely many such numbers where $k < K$.

As $\lim_{n \rightarrow \infty} x_n = c$, any $x \neq c$ appears at most finitely many times in $\{x_n\}_{n=1}^{\infty}$.

Hence, $\exists M$ such that for $n \geq M$, all the numbers x_n that are rational have a denominator larger than or equal to K .

$$\text{For } n \geq M, \quad |f(x_n) - 0| = f(x_n) \leq 1/K < \epsilon.$$

$\Rightarrow f$ is continuous at irrational c .

□

Example: Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) := 0$ if $x \neq 0$ and $g(0) := 1$.

Example: Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) := 0$ if $x \neq 0$ and $g(0) := 1$.

g is discontinuous at 0, but continuous on $\mathbb{R} \setminus \{0\}$.

Example: Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) := 0$ if $x \neq 0$ and $g(0) := 1$.

g is discontinuous at 0, but continuous on $\mathbb{R} \setminus \{0\}$.

$x = 0$ is called a *removable discontinuity*:

We could redefine $g(0)$ and obtain a continuous function.

Example: Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) := 0$ if $x \neq 0$ and $g(0) := 1$.

g is discontinuous at 0, but continuous on $\mathbb{R} \setminus \{0\}$.

$x = 0$ is called a *removable discontinuity*:

We could redefine $g(0)$ and obtain a continuous function.

The jump discontinuity, $f(x) := -1$ for $x < 0$ and $f(x) := 1$ for $x \geq 0$, does not have a removable singularity at $x = 0$.

Example: Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) := 0$ if $x \neq 0$ and $g(0) := 1$.

g is discontinuous at 0, but continuous on $\mathbb{R} \setminus \{0\}$.

$x = 0$ is called a *removable discontinuity*:

We could redefine $g(0)$ and obtain a continuous function.

The jump discontinuity, $f(x) := -1$ for $x < 0$ and $f(x) := 1$ for $x \geq 0$, does not have a removable singularity at $x = 0$.

$\lim_{x \rightarrow 0} g(x)$ exists $\lim_{x \rightarrow 0} f(x)$ does not.

Example: Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) := 0$ if $x \neq 0$ and $g(0) := 1$.

g is discontinuous at 0, but continuous on $\mathbb{R} \setminus \{0\}$.

$x = 0$ is called a *removable discontinuity*:

We could redefine $g(0)$ and obtain a continuous function.

The jump discontinuity, $f(x) := -1$ for $x < 0$ and $f(x) := 1$ for $x \geq 0$, does not have a removable singularity at $x = 0$.

$\lim_{x \rightarrow 0} g(x)$ exists $\lim_{x \rightarrow 0} f(x)$ does not.

Another thing:

Let $A := \{0\}$ and $B := \mathbb{R} \setminus \{0\}$.

Example: Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) := 0$ if $x \neq 0$ and $g(0) := 1$.

g is discontinuous at 0, but continuous on $\mathbb{R} \setminus \{0\}$.

$x = 0$ is called a *removable discontinuity*:

We could redefine $g(0)$ and obtain a continuous function.

The jump discontinuity, $f(x) := -1$ for $x < 0$ and $f(x) := 1$ for $x \geq 0$, does not have a removable singularity at $x = 0$.

$\lim_{x \rightarrow 0} g(x)$ exists $\lim_{x \rightarrow 0} f(x)$ does not.

Another thing:

Let $A := \{0\}$ and $B := \mathbb{R} \setminus \{0\}$.

$g|_A$ is continuous, but g is not continuous on A .

Example: Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) := 0$ if $x \neq 0$ and $g(0) := 1$.

g is discontinuous at 0, but continuous on $\mathbb{R} \setminus \{0\}$.

$x = 0$ is called a *removable discontinuity*:

We could redefine $g(0)$ and obtain a continuous function.

The jump discontinuity, $f(x) := -1$ for $x < 0$ and $f(x) := 1$ for $x \geq 0$, does not have a removable singularity at $x = 0$.

$\lim_{x \rightarrow 0} g(x)$ exists $\lim_{x \rightarrow 0} f(x)$ does not.

Another thing:

Let $A := \{0\}$ and $B := \mathbb{R} \setminus \{0\}$.

$g|_A$ is continuous, but g is not continuous on A .

$g|_B$ is continuous, and g is continuous on B .