

BA: 7.1

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

We wish to generalize convergence, continuity, etc.

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. *(nonnegativity)*

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. *(nonnegativity)*
- (ii) $d(x, y) = 0 \iff x = y$. *(identity of indiscernibles)*

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. *(nonnegativity)*
- (ii) $d(x, y) = 0 \iff x = y$. *(identity of indiscernibles)*
- (iii) $d(x, y) = d(y, x)$. *(symmetry)*

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. *(nonnegativity)*
- (ii) $d(x, y) = 0 \iff x = y$. *(identity of indiscernibles)*
- (iii) $d(x, y) = d(y, x)$. *(symmetry)*
- (iv) $d(x, z) \leq d(x, y) + d(y, z)$. *(triangle inequality)*

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. *(nonnegativity)*
- (ii) $d(x, y) = 0 \iff x = y$. *(identity of indiscernibles)*
- (iii) $d(x, y) = d(y, x)$. *(symmetry)*
- (iv) $d(x, z) \leq d(x, y) + d(y, z)$. *(triangle inequality)*

(X, d) is called a *metric space*.

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. (nonnegativity)
- (ii) $d(x, y) = 0 \iff x = y$. (identity of indiscernibles)
- (iii) $d(x, y) = d(y, x)$. (symmetry)
- (iv) $d(x, z) \leq d(x, y) + d(y, z)$. (triangle inequality)

(X, d) is called a *metric space*.

d is called the *metric* or the *distance function*.

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. (nonnegativity)
- (ii) $d(x, y) = 0 \iff x = y$. (identity of indiscernibles)
- (iii) $d(x, y) = d(y, x)$. (symmetry)
- (iv) $d(x, z) \leq d(x, y) + d(y, z)$. (triangle inequality)

(X, d) is called a *metric space*.

d is called the *metric* or the *distance function*.

Sometimes write just X instead of (X, d) if the metric is clear from context.

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. (nonnegativity)
- (ii) $d(x, y) = 0 \iff x = y$. (identity of indiscernibles)
- (iii) $d(x, y) = d(y, x)$. (symmetry)
- (iv) $d(x, z) \leq d(x, y) + d(y, z)$. (triangle inequality)

(X, d) is called a *metric space*.

d is called the *metric* or the *distance function*.

Sometimes write just X instead of (X, d) if the metric is clear from context.

Think of d as a distance between points.

We wish to generalize convergence, continuity, etc.

Definition

Let X be a set, and let $d: X \times X \rightarrow \mathbb{R}$ be such that for all $x, y, z \in X$

- (i) $d(x, y) \geq 0$. (nonnegativity)
- (ii) $d(x, y) = 0 \iff x = y$. (identity of indiscernibles)
- (iii) $d(x, y) = d(y, x)$. (symmetry)
- (iv) $d(x, z) \leq d(x, y) + d(y, z)$. (triangle inequality)

(X, d) is called a *metric space*.


d is called the *metric* or the *distance function*.

Sometimes write just X instead of (X, d) if the metric is clear from context.


Think of d as a distance between points.

(i)–(iii) have obvious geometric interpretation.

The interpretation of triangle inequality:



The interpretation of triangle inequality:



Warning: It is convenient to draw figures and diagrams in the plane, but that is just one particular metric space!

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.
(i)–(iii) are easy to verify.

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z)$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z|$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z|$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z|$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric* on \mathbb{R} .

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric* on \mathbb{R} . (It's the default one if none mentioned.)

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric* on \mathbb{R} . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} .

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric* on \mathbb{R} . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy.

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.


(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.



Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):


$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.

$d(x, z)$



Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):


$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.

$$d(x, z) = \varphi(|x - z|)$$



Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.

(i)–(iii) are easy to verify. For triangle inequality, (iv):


$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.

$$d(x, z) = \varphi(|x - z|) = \varphi(|x - y + y - z|)$$



Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.


(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.



$$d(x, z) = \varphi(|x - z|) = \varphi(|x - y + y - z|) \leq \varphi(|x - y| + |y - z|)$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.


(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.



$$d(x, z) = \varphi(|x - z|) = \varphi(|x - y + y - z|) \leq \varphi(|x - y| + |y - z|)$$

$$= \frac{|x - y| + |y - z|}{|x - y| + |y - z| + 1}$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.


(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.



$$d(x, z) = \varphi(|x - z|) = \varphi(|x - y + y - z|) \leq \varphi(|x - y| + |y - z|)$$

$$= \frac{|x - y| + |y - z|}{|x - y| + |y - z| + 1} = \frac{|x - y|}{|x - y| + |y - z| + 1} + \frac{|y - z|}{|x - y| + |y - z| + 1}$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.


(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.



$$d(x, z) = \varphi(|x - z|) = \varphi(|x - y + y - z|) \leq \varphi(|x - y| + |y - z|)$$

$$= \frac{|x - y| + |y - z|}{|x - y| + |y - z| + 1} = \frac{|x - y|}{|x - y| + |y - z| + 1} + \frac{|y - z|}{|x - y| + |y - z| + 1}$$

$$\leq \frac{|x - y|}{|x - y| + 1} + \frac{|y - z|}{|y - z| + 1}$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.


(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.



$$d(x, z) = \varphi(|x - z|) = \varphi(|x - y + y - z|) \leq \varphi(|x - y| + |y - z|)$$

$$= \frac{|x - y| + |y - z|}{|x - y| + |y - z| + 1} = \frac{|x - y|}{|x - y| + |y - z| + 1} + \frac{|y - z|}{|x - y| + |y - z| + 1}$$

$$\leq \frac{|x - y|}{|x - y| + 1} + \frac{|y - z|}{|y - z| + 1} = d(x, y) + d(y, z).$$

Example: \mathbb{R} is a metric space with the metric $d(x, y) := |x - y|$.


(i)–(iii) are easy to verify. For triangle inequality, (iv):

$$d(x, z) = |x - z| = |x - y + y - z| \leq |x - y| + |y - z| = d(x, y) + d(y, z).$$

d is the *standard metric on \mathbb{R}* . (It's the default one if none mentioned.)

Example: We can also have a different metric on \mathbb{R} . E.g., $d(x, y) := \frac{|x - y|}{|x - y| + 1}$.

(i)–(iii) are again easy. For triangle inequality, (iv), note $d(x, y) = \varphi(|x - y|)$, where $\varphi(t) = \frac{t}{t+1}$ and φ is increasing.



$$d(x, z) = \varphi(|x - z|) = \varphi(|x - y + y - z|) \leq \varphi(|x - y| + |y - z|)$$

$$= \frac{|x - y| + |y - z|}{|x - y| + |y - z| + 1} = \frac{|x - y|}{|x - y| + |y - z| + 1} + \frac{|y - z|}{|x - y| + |y - z| + 1}$$

$$\leq \frac{|x - y|}{|x - y| + 1} + \frac{|y - z|}{|y - z| + 1} = d(x, y) + d(y, z).$$

With this metric, $d(x, y) < 1$ for all $x, y \in \mathbb{R}$.

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$.

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. (We don't write \vec{x} nor \mathbf{x})

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. (We don't write \vec{x} nor \mathbf{x})

We also write $0 \in \mathbb{R}^n$ to mean $(0, 0, \dots, 0)$.

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. (We don't write \vec{x} nor \mathbf{x})

We also write $0 \in \mathbb{R}^n$ to mean $(0, 0, \dots, 0)$.

Lemma (Cauchy–Schwarz inequality)

If $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$, then

$$\left(\sum_{k=1}^n x_k y_k \right)^2 \leq \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right).$$

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. (We don't write \vec{x} nor \mathbf{x})

We also write $0 \in \mathbb{R}^n$ to mean $(0, 0, \dots, 0)$.

Lemma (Cauchy–Schwarz inequality)

If $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$, then

$$\left(\sum_{k=1}^n x_k y_k \right)^2 \leq \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right).$$

Proof: A square of a real number is nonnegative.

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. (We don't write \vec{x} nor \mathbf{x})

We also write $0 \in \mathbb{R}^n$ to mean $(0, 0, \dots, 0)$.

Lemma (Cauchy–Schwarz inequality)

If $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$, then

$$\left(\sum_{k=1}^n x_k y_k \right)^2 \leq \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right).$$

Proof: A square of a real number is nonnegative. Hence,

$$0 \leq \sum_{k=1}^n \sum_{\ell=1}^n (x_k y_\ell - x_\ell y_k)^2$$

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. (We don't write \vec{x} nor \mathbf{x})

We also write $0 \in \mathbb{R}^n$ to mean $(0, 0, \dots, 0)$.

Lemma (Cauchy–Schwarz inequality)

If $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$, then

$$\left(\sum_{k=1}^n x_k y_k \right)^2 \leq \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right).$$

Proof: A square of a real number is nonnegative. Hence,

$$0 \leq \sum_{k=1}^n \sum_{\ell=1}^n (x_k y_\ell - x_\ell y_k)^2 = \sum_{k=1}^n \sum_{\ell=1}^n (x_k^2 y_\ell^2 + x_\ell^2 y_k^2 - 2x_k x_\ell y_k y_\ell)$$

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. (We don't write \vec{x} nor \mathbf{x})

We also write $0 \in \mathbb{R}^n$ to mean $(0, 0, \dots, 0)$.

Lemma (Cauchy–Schwarz inequality)

If $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$, then

$$\left(\sum_{k=1}^n x_k y_k \right)^2 \leq \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right).$$

Proof: A square of a real number is nonnegative. Hence,

$$\begin{aligned} 0 &\leq \sum_{k=1}^n \sum_{\ell=1}^n (x_k y_\ell - x_\ell y_k)^2 = \sum_{k=1}^n \sum_{\ell=1}^n (x_k^2 y_\ell^2 + x_\ell^2 y_k^2 - 2x_k x_\ell y_k y_\ell) \\ &= \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{\ell=1}^n y_\ell^2 \right) + \left(\sum_{k=1}^n y_k^2 \right) \left(\sum_{\ell=1}^n x_\ell^2 \right) - 2 \left(\sum_{k=1}^n x_k y_k \right) \left(\sum_{\ell=1}^n x_\ell y_\ell \right). \end{aligned}$$

$\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}$ is the n -dimensional *euclidean space*.

Notation for points: $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. (We don't write \vec{x} nor \mathbf{x})

We also write $0 \in \mathbb{R}^n$ to mean $(0, 0, \dots, 0)$.

Lemma (Cauchy–Schwarz inequality)

If $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, $y = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n$, then

$$\left(\sum_{k=1}^n x_k y_k \right)^2 \leq \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right).$$

Proof: A square of a real number is nonnegative. Hence,

$$\begin{aligned} 0 &\leq \sum_{k=1}^n \sum_{\ell=1}^n (x_k y_\ell - x_\ell y_k)^2 = \sum_{k=1}^n \sum_{\ell=1}^n (x_k^2 y_\ell^2 + x_\ell^2 y_k^2 - 2x_k x_\ell y_k y_\ell) \\ &= \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{\ell=1}^n y_\ell^2 \right) + \left(\sum_{k=1}^n y_k^2 \right) \left(\sum_{\ell=1}^n x_\ell^2 \right) - 2 \left(\sum_{k=1}^n x_k y_k \right) \left(\sum_{\ell=1}^n x_\ell y_\ell \right). \\ \Rightarrow \quad 0 &\leq \left(\sum_{k=1}^n x_k^2 \right) \left(\sum_{k=1}^n y_k^2 \right) - \left(\sum_{k=1}^n x_k y_k \right)^2. \end{aligned}$$

□

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$(d(x, z))^2$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$(d(x, z))^2 = \sum_{k=1}^n (x_k - z_k)^2$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$(d(x, z))^2 = \sum_{k=1}^n (x_k - z_k)^2 = \sum_{k=1}^n (x_k - y_k + y_k - z_k)^2$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$\begin{aligned} (d(x, z))^2 &= \sum_{k=1}^n (x_k - z_k)^2 = \sum_{k=1}^n (x_k - y_k + y_k - z_k)^2 \\ &= \sum_{k=1}^n \left((x_k - y_k)^2 + (y_k - z_k)^2 + 2(x_k - y_k)(y_k - z_k) \right) \end{aligned}$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$\begin{aligned} (d(x, z))^2 &= \sum_{k=1}^n (x_k - z_k)^2 = \sum_{k=1}^n (x_k - y_k + y_k - z_k)^2 \\ &= \sum_{k=1}^n \left((x_k - y_k)^2 + (y_k - z_k)^2 + 2(x_k - y_k)(y_k - z_k) \right) \\ &= \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2 \sum_{k=1}^n (x_k - y_k)(y_k - z_k) \end{aligned}$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$\begin{aligned} (d(x, z))^2 &= \sum_{k=1}^n (x_k - z_k)^2 = \sum_{k=1}^n (x_k - y_k + y_k - z_k)^2 \\ &= \sum_{k=1}^n \left((x_k - y_k)^2 + (y_k - z_k)^2 + 2(x_k - y_k)(y_k - z_k) \right) \\ &= \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2 \sum_{k=1}^n (x_k - y_k)(y_k - z_k) \\ &\leq \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2\sqrt{\sum_{k=1}^n (x_k - y_k)^2 \sum_{k=1}^n (y_k - z_k)^2} \end{aligned}$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$\begin{aligned} (d(x, z))^2 &= \sum_{k=1}^n (x_k - z_k)^2 = \sum_{k=1}^n (x_k - y_k + y_k - z_k)^2 \\ &= \sum_{k=1}^n \left((x_k - y_k)^2 + (y_k - z_k)^2 + 2(x_k - y_k)(y_k - z_k) \right) \\ &= \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2 \sum_{k=1}^n (x_k - y_k)(y_k - z_k) \\ &\leq \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2\sqrt{\sum_{k=1}^n (x_k - y_k)^2 \sum_{k=1}^n (y_k - z_k)^2} \\ &= \left(\sqrt{\sum_{k=1}^n (x_k - y_k)^2} + \sqrt{\sum_{k=1}^n (y_k - z_k)^2} \right)^2 \end{aligned}$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$\begin{aligned} (d(x, z))^2 &= \sum_{k=1}^n (x_k - z_k)^2 = \sum_{k=1}^n (x_k - y_k + y_k - z_k)^2 \\ &= \sum_{k=1}^n \left((x_k - y_k)^2 + (y_k - z_k)^2 + 2(x_k - y_k)(y_k - z_k) \right) \\ &= \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2 \sum_{k=1}^n (x_k - y_k)(y_k - z_k) \\ &\leq \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2\sqrt{\sum_{k=1}^n (x_k - y_k)^2 \sum_{k=1}^n (y_k - z_k)^2} \\ &= \left(\sqrt{\sum_{k=1}^n (x_k - y_k)^2} + \sqrt{\sum_{k=1}^n (y_k - z_k)^2} \right)^2 = (d(x, y) + d(y, z))^2. \end{aligned}$$

Example: We define the standard metric for \mathbb{R}^n :

$$d(x, y) := \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_n - y_n)^2} = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

Note: If $n = 1$ agrees with above.

The only tricky bit (again) is to check triangle inequality.

$$\begin{aligned} (d(x, z))^2 &= \sum_{k=1}^n (x_k - z_k)^2 = \sum_{k=1}^n (x_k - y_k + y_k - z_k)^2 \\ &= \sum_{k=1}^n \left((x_k - y_k)^2 + (y_k - z_k)^2 + 2(x_k - y_k)(y_k - z_k) \right) \\ &= \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2 \sum_{k=1}^n (x_k - y_k)(y_k - z_k) \\ &\leq \sum_{k=1}^n (x_k - y_k)^2 + \sum_{k=1}^n (y_k - z_k)^2 + 2\sqrt{\sum_{k=1}^n (x_k - y_k)^2 \sum_{k=1}^n (y_k - z_k)^2} \\ &= \left(\sqrt{\sum_{k=1}^n (x_k - y_k)^2} + \sqrt{\sum_{k=1}^n (y_k - z_k)^2} \right)^2 = (d(x, y) + d(y, z))^2. \end{aligned}$$

The triangle inequality follows as $\sqrt{\cdot}$ is an increasing function.

Example: Complex numbers \mathbb{C} is the set of numbers $z = x + iy$, where $x, y \in \mathbb{R}$.

Example: Complex numbers \mathbb{C} is the set of numbers $z = x + iy$, where $x, y \in \mathbb{R}$.

Imposing $i^2 = -1$, makes \mathbb{C} into a field.

Example: Complex numbers \mathbb{C} is the set of numbers $z = x + iy$, where $x, y \in \mathbb{R}$.

Imposing $i^2 = -1$, makes \mathbb{C} into a field.

To make \mathbb{C} into a metric space, identify \mathbb{C} with \mathbb{R}^2 by

$$x + iy \in \mathbb{C} \quad \leftrightarrow \quad (x, y) \in \mathbb{R}^2,$$

and use the standard metric.

Example: Complex numbers \mathbb{C} is the set of numbers $z = x + iy$, where $x, y \in \mathbb{R}$.

Imposing $i^2 = -1$, makes \mathbb{C} into a field.

To make \mathbb{C} into a metric space, identify \mathbb{C} with \mathbb{R}^2 by

$$x + iy \in \mathbb{C} \quad \leftrightarrow \quad (x, y) \in \mathbb{R}^2,$$

and use the standard metric.

Define the *complex modulus* $|x + iy| := \sqrt{x^2 + y^2}$.

Example: Complex numbers \mathbb{C} is the set of numbers $z = x + iy$, where $x, y \in \mathbb{R}$.

Imposing $i^2 = -1$, makes \mathbb{C} into a field.

To make \mathbb{C} into a metric space, identify \mathbb{C} with \mathbb{R}^2 by

$$x + iy \in \mathbb{C} \quad \leftrightarrow \quad (x, y) \in \mathbb{R}^2,$$

and use the standard metric.

Define the *complex modulus* $|x + iy| := \sqrt{x^2 + y^2}$.

If $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$, then

$$d(z_1, z_2) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} = |z_1 - z_2|.$$

Example: For any set X , define the *discrete metric*:

$$d(x, y) := \begin{cases} 1 & \text{if } x \neq y, \\ 0 & \text{if } x = y. \end{cases}$$

Example: For any set X , define the *discrete metric*:

$$d(x, y) := \begin{cases} 1 & \text{if } x \neq y, \\ 0 & \text{if } x = y. \end{cases}$$


Easy to check the properties.

Example: For any set X , define the *discrete metric*:

$$d(x, y) := \begin{cases} 1 & \text{if } x \neq y, \\ 0 & \text{if } x = y. \end{cases}$$

Easy to check the properties.

E.g., if $X = \{a, b, c, d, e\}$, then




Example: For any set X , define the *discrete metric*:

$$d(x, y) := \begin{cases} 1 & \text{if } x \neq y, \\ 0 & \text{if } x = y. \end{cases}$$

Easy to check the properties.

E.g., if $X = \{a, b, c, d, e\}$, then



A very useful “smell test” for statements about metric spaces.

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$.

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as

$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0.$
 $d(f, g) = 0$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as $d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|$.

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x)$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as $d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|$.

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x) \Rightarrow f = g$.

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.

$d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x) \Rightarrow f = g$.

(iii) $d(f, g) = d(g, f)$ is trivial.

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as $d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|$.

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x) \Rightarrow f = g$.

(iii) $d(f, g) = d(g, f)$ is trivial.

(iv) $d(f, g)$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x) \Rightarrow f = g$.

(iii) $d(f, g) = d(g, f)$ is trivial.

(iv) $d(f, g) = \sup_{x \in [a, b]} |f(x) - g(x)|$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x) \Rightarrow f = g$.

(iii) $d(f, g) = d(g, f)$ is trivial.

(iv) $d(f, g) = \sup_{x \in [a, b]} |f(x) - g(x)| = \sup_{x \in [a, b]} |f(x) - h(x) + h(x) - g(x)|$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as $d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|$.

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x) \Rightarrow f = g$.

(iii) $d(f, g) = d(g, f)$ is trivial.

(iv) $d(f, g) = \sup_{x \in [a, b]} |f(x) - g(x)| = \sup_{x \in [a, b]} |f(x) - h(x) + h(x) - g(x)|$
 $\leq \sup_{x \in [a, b]} (|f(x) - h(x)| + |h(x) - g(x)|)$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x) \Rightarrow f = g$.

(iii) $d(f, g) = d(g, f)$ is trivial.

(iv)
$$\begin{aligned} d(f, g) &= \sup_{x \in [a, b]} |f(x) - g(x)| = \sup_{x \in [a, b]} |f(x) - h(x) + h(x) - g(x)| \\ &\leq \sup_{x \in [a, b]} (|f(x) - h(x)| + |h(x) - g(x)|) \\ &\leq \sup_{x \in [a, b]} |f(x) - h(x)| + \sup_{x \in [a, b]} |h(x) - g(x)| \end{aligned}$$

Example: Let $C([a, b], \mathbb{R})$ be the set of continuous functions $f: [a, b] \rightarrow \mathbb{R}$. Define the (standard) metric as
$$d(f, g) := \sup_{x \in [a, b]} |f(x) - g(x)|.$$

$d(f, g)$ is finite as $|f(x) - g(x)|$ is a continuous function on a closed bounded interval $[a, b]$ and so bounded.

(i) Clearly $d(f, g) \geq 0$.

(ii) $f = g \Rightarrow \forall x, |f(x) - g(x)| = 0 \Rightarrow d(f, g) = 0$.
 $d(f, g) = 0 \Rightarrow \forall x, |f(x) - g(x)| \leq d(f, g) = 0 \Rightarrow \forall x, f(x) = g(x) \Rightarrow f = g$.


(iii) $d(f, g) = d(g, f)$ is trivial.

(iv)
$$\begin{aligned} d(f, g) &= \sup_{x \in [a, b]} |f(x) - g(x)| = \sup_{x \in [a, b]} |f(x) - h(x) + h(x) - g(x)| \\ &\leq \sup_{x \in [a, b]} (|f(x) - h(x)| + |h(x) - g(x)|) \\ &\leq \sup_{x \in [a, b]} |f(x) - h(x)| + \sup_{x \in [a, b]} |h(x) - g(x)| = d(f, h) + d(h, g). \end{aligned}$$

Example (Great circle distance): $S^2 = \{x \in \mathbb{R}^3 : x_1^2 + x_2^2 + x_3^2 = 1\}$ is the unit sphere in \mathbb{R}^3 .

Example (Great circle distance): $S^2 = \{x \in \mathbb{R}^3 : x_1^2 + x_2^2 + x_3^2 = 1\}$ is the unit sphere in \mathbb{R}^3 .

If x and y are in S^2 , the lines they make with the origin meet at angle θ (in radians).




Example (Great circle distance): $S^2 = \{x \in \mathbb{R}^3 : x_1^2 + x_2^2 + x_3^2 = 1\}$ is the unit sphere in \mathbb{R}^3 .

If x and y are in S^2 , the lines they make with the origin meet at angle θ (in radians).

The function defined by

$$d(x, y) := \theta$$




Example (Great circle distance): $S^2 = \{x \in \mathbb{R}^3 : x_1^2 + x_2^2 + x_3^2 = 1\}$ is the unit sphere in \mathbb{R}^3 .

If x and y are in S^2 , the lines they make with the origin meet at angle θ (in radians).

The function defined by

$d(x, y) := \theta = \arccos(x_1y_1 + x_2y_2 + x_3y_3)$
gives a metric (we skip the proof).




Example (Great circle distance): $S^2 = \{x \in \mathbb{R}^3 : x_1^2 + x_2^2 + x_3^2 = 1\}$ is the unit sphere in \mathbb{R}^3 .

If x and y are in S^2 , the lines they make with the origin meet at angle θ (in radians).

The function defined by

$d(x, y) := \theta = \arccos(x_1y_1 + x_2y_2 + x_3y_3)$
gives a metric (we skip the proof).

It is the shortest path between the points if we travel along the sphere.



Proposition

Let (X, d) be a metric space and $Y \subset X$. Then the restriction $d|_{Y \times Y}$ is a metric on Y .

Proposition

Let (X, d) be a metric space and $Y \subset X$. Then the restriction $d|_{Y \times Y}$ is a metric on Y .

Proof: Obvious.

Proposition

Let (X, d) be a metric space and $Y \subset X$. Then the restriction $d|_{Y \times Y}$ is a metric on Y .

Proof: Obvious.

Definition

If (X, d) is a metric space, $Y \subset X$, and $d' := d|_{Y \times Y}$, then (Y, d') is a *subspace* of (X, d) , and d' is the *subspace metric* (sometimes we talk about *subspace topology*).

Proposition

Let (X, d) be a metric space and $Y \subset X$. Then the restriction $d|_{Y \times Y}$ is a metric on Y .

Proof: Obvious.

Definition

If (X, d) is a metric space, $Y \subset X$, and $d' := d|_{Y \times Y}$, then (Y, d') is a *subspace* of (X, d) , and d' is the *subspace metric* (sometimes we talk about *subspace topology*).

Common to just write d for the metric on Y .

Definition

Let (X, d) be a metric space. $S \subset X$ is *bounded* if $\exists p \in X$ and $B \in \mathbb{R}$ such that

$$d(p, x) \leq B \quad \text{for all } x \in S.$$

Definition

Let (X, d) be a metric space. $S \subset X$ is *bounded* if $\exists p \in X$ and $B \in \mathbb{R}$ such that

$$d(p, x) \leq B \quad \text{for all } x \in S.$$

(X, d) is bounded if X is a bounded subset.

Definition

Let (X, d) be a metric space. $S \subset X$ is *bounded* if $\exists p \in X$ and $B \in \mathbb{R}$ such that

$$d(p, x) \leq B \quad \text{for all } x \in S.$$

(X, d) is bounded if X is a bounded subset.

Example: \mathbb{R} with standard metric is not bounded.

Definition

Let (X, d) be a metric space. $S \subset X$ is *bounded* if $\exists p \in X$ and $B \in \mathbb{R}$ such that

$$d(p, x) \leq B \quad \text{for all } x \in S.$$

(X, d) is bounded if X is a bounded subset.

Example: \mathbb{R} with standard metric is not bounded.

Example: \mathbb{R} with the discrete metric is bounded.

Definition

Let (X, d) be a metric space. $S \subset X$ is *bounded* if $\exists p \in X$ and $B \in \mathbb{R}$ such that

$$d(p, x) \leq B \quad \text{for all } x \in S.$$

(X, d) is bounded if X is a bounded subset.

Example: \mathbb{R} with standard metric is not bounded.

Example: \mathbb{R} with the discrete metric is bounded.

Exercise: Suppose $X \neq \emptyset$. Then $S \subset X$ being bounded is equivalent to

Definition

Let (X, d) be a metric space. $S \subset X$ is *bounded* if $\exists p \in X$ and $B \in \mathbb{R}$ such that

$$d(p, x) \leq B \quad \text{for all } x \in S.$$

(X, d) is bounded if X is a bounded subset.

Example: \mathbb{R} with standard metric is not bounded.

Example: \mathbb{R} with the discrete metric is bounded.

Exercise: Suppose $X \neq \emptyset$. Then $S \subset X$ being bounded is equivalent to

(i) $\forall p \in X, \exists B > 0$ such that $d(p, x) \leq B$ for all $x \in S$.

Definition

Let (X, d) be a metric space. $S \subset X$ is *bounded* if $\exists p \in X$ and $B \in \mathbb{R}$ such that

$$d(p, x) \leq B \quad \text{for all } x \in S.$$

(X, d) is bounded if X is a bounded subset.

Example: \mathbb{R} with standard metric is not bounded.

Example: \mathbb{R} with the discrete metric is bounded.

Exercise: Suppose $X \neq \emptyset$. Then $S \subset X$ being bounded is equivalent to

- (i) $\forall p \in X, \exists B > 0$ such that $d(p, x) \leq B$ for all $x \in S$.
- (ii) $\text{diam}(S) := \sup\{d(x, y) : x, y \in S\} < \infty$.

Definition

Let (X, d) be a metric space. $S \subset X$ is *bounded* if $\exists p \in X$ and $B \in \mathbb{R}$ such that

$$d(p, x) \leq B \quad \text{for all } x \in S.$$

(X, d) is bounded if X is a bounded subset.

Example: \mathbb{R} with standard metric is not bounded.

Example: \mathbb{R} with the discrete metric is bounded.

Exercise: Suppose $X \neq \emptyset$. Then $S \subset X$ being bounded is equivalent to

- (i) $\forall p \in X, \exists B > 0$ such that $d(p, x) \leq B$ for all $x \in S$.
- (ii) $\text{diam}(S) := \sup\{d(x, y) : x, y \in S\} < \infty$.

The quantity $\text{diam}(S)$ is called the *diameter* of a set.

Exercise: Let (X, d_X) and (Y, d_Y) be metric spaces.

- a) Show that $(X \times Y, d)$ with $d((x_1, y_1), (x_2, y_2)) := d_X(x_1, x_2) + d_Y(y_1, y_2)$ is a metric space.
- b) Show that $(X \times Y, d)$ with $d((x_1, y_1), (x_2, y_2)) := \max\{d_X(x_1, x_2), d_Y(y_1, y_2)\}$ is a metric space.

Exercise: Let (X, d_X) and (Y, d_Y) be metric spaces.

- a) Show that $(X \times Y, d)$ with $d((x_1, y_1), (x_2, y_2)) := d_X(x_1, x_2) + d_Y(y_1, y_2)$ is a metric space.
- b) Show that $(X \times Y, d)$ with $d((x_1, y_1), (x_2, y_2)) := \max\{d_X(x_1, x_2), d_Y(y_1, y_2)\}$ is a metric space.

Exercise: Let X be the set of continuous functions on $[0, 1]$. Let $\varphi: [0, 1] \rightarrow (0, \infty)$ be continuous. Define

$$d(f, g) := \int_0^1 |f(x) - g(x)| \varphi(x) dx.$$

Show that (X, d) is a metric space.